

Research Article

Application of Clifford Algebra on Group Theory

Farooqhusain Inamdar^{*}, Hasan S. N.

Department of Mathematics, Maulana Azad National Urdu University, Hyderabad, India
E-mail: farooqhusain2690@gmail.com

Received: 14 November 2023; **Revised:** 18 December 2023; **Accepted:** 18 March 2024

Abstract: The orthogonal operators defined as similarity transformations on Euclidean space E can also be considered as group actions on the Clifford Algebra. In this paper, we investigate the finite subgroup of Euclidean space E of Geometric Algebra over a finite dimension vector space E . The hierarchy of the finite subgroups of Clifford Algebra $C(E)$ is depicted through the lattice structure and we discussed the group action of these subgroups on the vector space E . Further, we shall address the number of non-trivial finite subgroups, Normal subgroups, and subnormal series of the subgroup B_3 of Clifford Algebra $C(E)$ constructed over the vector space E by performing group action $\Psi : B_3 \times B_3 \rightarrow B_3$ over the subgroup B_3 of Clifford Algebra $C(E)$.

Keywords: Geometric Algebra, Clifford Algebra, group action, equivalence class, principle homogeneous space, subnormal series, solvable group

MSC: 03C05, 03E20

1. Introduction

William K. Clifford introduced the Geometric Algebra which is now usually known as Clifford Algebra [1] by creating a new multiplication rule in Grassmann's Exterior Algebra [2, 3]. Clifford was able to unite two seemingly unrelated mathematical frameworks: the algebra of extensions, which Hermann Grassmann invented, and the quaternion, which Sir William Rowan Hamilton constructed [4]. Clifford observed that Hamilton and Grassmann were tackling the same problem from different perspectives, and consequently, he combined both ideas to define the Geometric Product in 1876. In 1878, Clifford made a significant contribution by integrating the framework of quaternion into Grassmann's algebra of extensions [5]. This combination resulted in a system specifically designed to align with the orthogonal geometry of any given space. Within this system, quaternion emerged as a specific instance of a broader category known as Clifford algebra. However, his ideas were not further developed by his contemporary mathematicians for many years [6].

The Clifford algebras have been developed with the involvement of several Mathematicians and physicists such as Rudolf Lipschitz, Theodor Valen, Elie Cartan, Claude Chevalley reinvented the "Clifford Algebra" [7] and established its power as a formal mathematics and physics language. Specifically, David Heston and Elie Cartan are notable contributors to the progress and development of Clifford algebra. Elie Cartan presented the idea of the spinor in 1913 and in 1938 the idea of the pure spinor and he defined Clifford algebra's as algebras of matrices and found that 8 has a periodicity inside

these algebraic structures, for more info, refer [8]. David Heston extended the concept of “Clifford Algebra” to devise a formalism and calls it Geometric Algebra [6]. He defines orthogonal operators as similarity transformations on Euclidian space E , which can also be considered as group actions in Clifford Algebra on the underlying Vector Space. In this paper, we investigate the finite subgroup of Euclidian space of Clifford Algebra over a finite dimension vector space E and we will show that the subgroup B_3 of Clifford algebra is Solvable.

2. Preliminaries

2.1 Clifford Algebra

The algebra denoted as $C(E)$, which is defined over an n -dimensional vector space E , includes a Clifford map denoted as $\rho : E \rightarrow C(E)$. This Clifford map must fulfil the condition that for any algebra L and any other Clifford map $\psi : E \rightarrow L$, there exists a unique algebra homomorphism $\psi_* : C(E) \rightarrow L$ see the Figure 1, such that $\psi_* \circ \rho = \psi$ [5, 7].




Figure 1. Unique algebra homomorphism

2.2 Clifford’s original definition

Grassmann’s exterior algebra $\wedge R^n$ of the linear space R^n is an associative algebra of dimension 2^n . In terms of a basis $\{\sigma_1, \sigma_2, \sigma_3, \dots, \sigma_n\}$ for R^n the exterior algebra $\wedge R^n$ has a basis,

$$1$$

$$\sigma_1, \sigma_2, \sigma_3, \dots, \sigma_n$$

$$\sigma_1 \wedge \sigma_2, \sigma_1 \wedge \sigma_3, \dots, \sigma_1 \wedge \sigma_n, \sigma_2 \wedge \sigma_3, \dots, \sigma_{n-1} \wedge \sigma_n$$

.

.

.

$$\sigma_1 \wedge \sigma_2 \wedge \sigma_3 \dots \wedge \sigma_n$$

The exterior algebra has unit 1 and satisfies the multiplication rules

$$\begin{aligned} \sigma_i \wedge \sigma_j &= -\sigma_j \wedge \sigma_i \text{ for } i \neq j \\ \sigma_i \wedge \sigma_i &= 0 \end{aligned} \tag{1}$$

Clifford in 1882 kept the first rule but altered the second rule, and arrived at the multiplication rule

$$\begin{aligned} \sigma_i \sigma_j &= -\sigma_j \sigma_i \text{ for } i \neq j \\ \sigma_i \sigma_i &= 1 \end{aligned} \tag{2}$$

This time $\{\sigma_1, \sigma_2, \sigma_3, \dots, \sigma_n\}$ is an orthonormal basis for the positive definite Euclidean space R^n . An associative algebra of dimension 2^n so defined is the Clifford algebra Cl_n .

Clifford in 1878, considered the multiplication rules

$$\begin{aligned} \sigma_i \sigma_j &= -\sigma_j \sigma_i \text{ for } i \neq j \\ \sigma_i \sigma_i &= -1 \end{aligned} \tag{3}$$

of the Clifford algebra $Cl_{(0, n)}$ of the negative definite space $R^{(0, n)}$ [7, 9].

2.3 Clifford mapping

A linear mapping $\psi : E \rightarrow L$ is referred to as a Clifford map from E to L provided that it meets the following criteria:

$$[\psi(x)]^2 = g(x, x)1_L = \|x\|^2 1_L \tag{4}$$

Which is equivalent to,

$$\psi(x)\psi(y) + \psi(y)\psi(x) = 2g(x, y)1_L \quad \forall x, y \in E \tag{5}$$

2.4 Even Clifford Algebra

Assume that E is a vector space having n dimensions and $\{\sigma_i\}_{i=1}^n$ is a basis of E then the 2^n elements $\sigma_{i_1}, \sigma_{i_2}, \dots, \sigma_{i_s}$ for $i_1 < i_2 < \dots < i_s, 0 \leq s \leq n$ form a basis for the Clifford Algebra $C(E)$. Let $L = C(E)$ and $\psi : E \rightarrow C(E)$ be given

by $\psi(u) = -u$ then we find a isomorphism ψ_* of $C(E)$ into itself so that $\psi_*(u) = -u$, $u \in E$, and may be written as $\psi_*(a) = a'$.

Now we put,

$$C^+(E) = \{a \in C(E) : a' = a\} \ \& \ C^-(E) = \{a \in C(E) : a' = -a\}, \ \text{also } \sigma'_s = (-1)^{|s|} \sigma_s.$$

Therefore the product $\sigma_1 \sigma_2 \sigma_3 \dots \sigma_s$ belongs to $C^+(E)$ or $C^-(E)$ according as 's' is even or odd, then the elements $\sigma_{i_1} \sigma_{i_2} \dots \sigma_{i_s}$ for $i_1 < i_2 < \dots < i_s$, $0 \leq s \leq n$ with even or odd s forms a basis of $C^+(E)$ or $C^-(E)$ respectively. Thus, Clifford Algebra is the direct sum of even part $C^+(E)$ and odd $C^-(E)$, i.e,

$$C(E) = C^+(E) \oplus C^-(E) \tag{6}$$

The even part is not only a subspace but also a sub-algebra of $C(E)$ which we call Even Clifford Algebra [5, 7, 10].

2.5 Arbitrary element of Clifford Algebra

The following formal polynomial represents an arbitrary element \mathcal{A} in the Clifford Algebra $Cl_{(p, q)}$:

$$\begin{aligned} \mathcal{A} = & a^0 \sigma_0 + \sum_{i=1}^n a^i \sigma_i + \sum_{i=1}^n \sum_{j=1}^n a^{ij} \sigma_{ij} + \dots + \sum_{i_1=1}^n \dots \sum_{i_k=1}^n a^{i_1 \dots i_k} \sigma_{i_1 \dots i_k} \\ & + \dots + a^{12 \dots n} \sigma_{12 \dots n} = \sum_{k=0}^n a^{i_1 i_2 \dots i_k} \sigma_{i_1 i_2 \dots i_k} \end{aligned} \tag{7}$$

2.6 Fundamental automorphism of Clifford Algebra

Clifford Algebra $Cl_{(p, q)}$ has four fundamental automorphism, which are as follows [7, 11, 12]:

2.6.1 Identity

Let \mathcal{A} be any random element of Clifford Algebra $Cl_{(p, q)}$, the Identity automorphism from $\mathcal{A} \rightarrow \mathcal{A}$ is one which carries $\sigma_i \rightarrow \sigma_i$.

2.6.2 Involution

Let $\mathcal{A} = \mathcal{A}' + \mathcal{A}''$ be the decomposition of an element of Clifford Algebra $Cl_{(p, q)}$, where \mathcal{A}' and \mathcal{A}'' contains homogeneous odd and even components individually, then automorphism $\mathcal{A} \rightarrow \mathcal{A}^*$ is the Involution so that the sign of the elements of \mathcal{A}'' doesn't change and the sign of elements of \mathcal{A}' changes, i.e,

$$\mathcal{A}^* = -\mathcal{A}' + \mathcal{A}''$$

In general Involution automorphism carries $\sigma_i \rightarrow -\sigma_i$, for any element of \mathcal{A} .

The Involution automorphism can also be expressed with the help of the volume element ω , i.e, $\omega = \sigma_{i_1 i_2 \dots i_{p+q}}$, such that, $\mathcal{A} = \omega \mathcal{A} \omega^{-1}$, where $\omega^{-1} = (-1)^{\frac{(p+q)(p+q-1)}{2}} \omega$ [1, 11].

2.6.3 Reversion

The Reversion of any element of Clifford Algebra $Cl_{(p, q)}$ is the Antiautomorphism from $\mathcal{A} \rightarrow \widetilde{\mathcal{A}}$, that is an alternative to any basis element $\sigma_{i_1 i_2 \dots i_k} \in \mathcal{A}$ by an element of $\sigma_{i_k i_{k-1} \dots i_1}$, such that:

$$\sigma_{i_1 i_2 \dots i_k} = (-1)^{\frac{k(k-1)}{2}} \sigma_{i_k i_{k-1} \dots i_1}$$

Hence for any element \mathcal{A} of Clifford Algebra $Cl_{(p, q)}$,

$$\widetilde{\mathcal{A}} = (-1)^{\frac{k(k-1)}{2}} \mathcal{A}$$

2.6.4 Conjugation

The Conjugation of any element \mathcal{A} of Clifford Algebra $Cl_{(p, q)}$ is the Antiautomorphism from $\mathcal{A} \rightarrow \widetilde{\mathcal{A}}^*$ which is the composition of Involution and Reversion Automorphism, [11] such that

$$\widetilde{\mathcal{A}}^* = (-1)^{\frac{k(k-1)}{2}} \mathcal{A}$$

3. The lattice structure of finite sub-group of Clifford Algebra




Figure 2. Lattice structure of Clifford Algebra

In this section, we give the classification of finite sub-groups of Clifford Algebra. The hierarchy of sub groups is represented in the lattice structure format which is given in the Figure 2.

4. Group action

Let G be a set of any elements and a binary operation $(*)$ together satisfied the Closure law, Associative law, and if there exist Unit element and Inverse of that element then we say that the set G is a group under the operation $(*)$. If the above group G also satisfied the commutative law then we called it as Abelian group [13].

Similarly, we can easily define the Subgroup, Normal group, Permutation group etc, for clarity refer [13, 14]. Here our main aim is to explain the Group Action which we will explain below.

Cayley's theorem states that it is possible to find a subgroup of the Permutations group $S(X)$, which is isomorphic to any given group. Let G be a subgroup of $S(X)$. Then, for any element $a \in G$ and any element $x \in X$, there exists an element $a(x) \in X$, and this relationship satisfies certain conditions.

- $e(x) = x$, where e embodies the identity in G .
- $(a, b)(x) = a(b(x)), \forall x \in X \ \& \ a, b \in G$.

The definition that follows abstracts this.

If X is any non-empty set, then let G be a group. Any mapping $\theta : G \times X \rightarrow X$ that satisfies the following criteria is an action of G on X [15].

1. $\theta(e, x) = x \ \forall x \in X$, where e embodies the identity in G .
2. $\theta(ab, x) = \theta(a, \theta(b, x)) \ \forall a, b \in G$.

If there exists a situation where G performs an action on X , it can be said that G acts on X . We study the finite groups and their action on various subgroup E of $C(E)$ and their subgroups which in turn are subgroups of $C(E)$.

Let us consider the following finite subgroup with respect to the operation geometric product.

$$B_3 = \{\pm 1, \pm\sigma_1, \pm\sigma_2, \pm\sigma_3, \pm\sigma_1\sigma_2, \pm\sigma_1\sigma_3, \pm\sigma_2\sigma_3, \pm\sigma_1\sigma_2\sigma_3\} \quad (8)$$

The composition table given in the Table 1 shows the action of B_3 on itself defined by: $\varphi : B_3 \times B_3 \rightarrow B_3$ so that $\varphi(x, y) = xy \ \forall x, y \in B_3$ and the subgroup is,

$$B_3 = \pm 1, \pm\sigma_1, \pm\sigma_2, \pm\sigma_1\sigma_2 \quad (9)$$

It can be verified that φ represents a group action on B_3 .

Table 1. Composition table

	1	-1	σ_1	$-\sigma_1$	σ_2	$-\sigma_2$	σ_3	$-\sigma_3$	$\sigma_1\sigma_2$	$\sigma_2\sigma_1$	$\sigma_1\sigma_3$	$\sigma_3\sigma_1$	$\sigma_2\sigma_3$	$\sigma_3\sigma_2$	$\sigma_1\sigma_2\sigma_3$	$\sigma_3\sigma_2\sigma_1$
1	1	-1	σ_1	$-\sigma_1$	σ_2	$-\sigma_2$	σ_3	$-\sigma_3$	$\sigma_1\sigma_2$	$\sigma_2\sigma_1$	$\sigma_1\sigma_3$	$\sigma_3\sigma_1$	$\sigma_2\sigma_3$	$\sigma_3\sigma_2$	$\sigma_1\sigma_2\sigma_3$	$\sigma_3\sigma_2\sigma_1$
-1	-1	1	$-\sigma_1$	σ_1	$-\sigma_2$	σ_2	σ_3	$-\sigma_3$	$\sigma_1\sigma_2$	$\sigma_2\sigma_1$	$\sigma_1\sigma_3$	$\sigma_3\sigma_1$	$\sigma_2\sigma_3$	$\sigma_3\sigma_2$	$\sigma_1\sigma_2\sigma_3$	$\sigma_3\sigma_2\sigma_1$
σ_1	σ_1	$-\sigma_1$	1	-1	$\sigma_1\sigma_2$	$\sigma_2\sigma_1$	$\sigma_1\sigma_3$	$\sigma_3\sigma_1$	σ_2	$-\sigma_2$	$\sigma_1\sigma_2\sigma_3$	$\sigma_3\sigma_2\sigma_1$	$-\sigma_3$	σ_3	$\sigma_2\sigma_3$	$\sigma_3\sigma_2$
$-\sigma_1$	$-\sigma_1$	σ_1	-1	1	$\sigma_2\sigma_1$	$\sigma_1\sigma_2$	$\sigma_3\sigma_1$	$\sigma_1\sigma_3$	$-\sigma_2$	σ_2	$\sigma_3\sigma_2\sigma_1$	$\sigma_1\sigma_2\sigma_3$	σ_3	$-\sigma_3$	$\sigma_3\sigma_2$	$\sigma_2\sigma_3$
σ_2	σ_2	$-\sigma_2$	$\sigma_2\sigma_1$	$\sigma_1\sigma_2$	1	-1	$\sigma_2\sigma_3$	$\sigma_3\sigma_2$	$-\sigma_1$	σ_1	σ_3	$-\sigma_3$	$\sigma_1\sigma_2\sigma_3$	$\sigma_3\sigma_2\sigma_1$	$\sigma_3\sigma_1$	$\sigma_1\sigma_3$
$-\sigma_2$	$-\sigma_2$	σ_2	$\sigma_1\sigma_2$	$\sigma_2\sigma_1$	-1	1	$\sigma_3\sigma_2$	$\sigma_2\sigma_3$	σ_1	$-\sigma_1$	$-\sigma_3$	σ_3	$\sigma_3\sigma_2\sigma_1$	$\sigma_1\sigma_2\sigma_3$	$\sigma_1\sigma_3$	$\sigma_3\sigma_1$
σ_3	σ_3	$-\sigma_3$	$\sigma_3\sigma_1$	$\sigma_1\sigma_3$	$\sigma_3\sigma_2$	$\sigma_2\sigma_3$	1	-1	$\sigma_1\sigma_2\sigma_3$	$\sigma_3\sigma_2\sigma_1$	$-\sigma_2$	σ_2	σ_1	$-\sigma_1$	$\sigma_1\sigma_2$	$\sigma_2\sigma_1$
$-\sigma_3$	$-\sigma_3$	σ_3	$\sigma_1\sigma_3$	$\sigma_3\sigma_1$	$\sigma_2\sigma_3$	$\sigma_3\sigma_2$	-1	1	$\sigma_3\sigma_2\sigma_1$	$\sigma_1\sigma_2\sigma_3$	σ_2	$-\sigma_2$	$-\sigma_1$	σ_1	$\sigma_2\sigma_1$	$\sigma_1\sigma_2$
$\sigma_1\sigma_2$	$\sigma_1\sigma_2$	$\sigma_2\sigma_1$	$-\sigma_2$	σ_2	σ_1	$-\sigma_1$	$\sigma_1\sigma_2\sigma_3$	$\sigma_3\sigma_2\sigma_1$	-1	1	$\sigma_1\sigma_3$	$\sigma_3\sigma_1$	$\sigma_2\sigma_3$	$\sigma_3\sigma_2$	$-\sigma_3$	σ_3
$\sigma_2\sigma_1$	$\sigma_2\sigma_1$	$\sigma_1\sigma_2$	σ_2	$-\sigma_2$	$-\sigma_1$	σ_1	$\sigma_3\sigma_2\sigma_1$	$\sigma_1\sigma_2\sigma_3$	1	-1	$\sigma_3\sigma_1$	$\sigma_1\sigma_3$	$\sigma_3\sigma_2$	$\sigma_2\sigma_3$	σ_3	$-\sigma_3$
$\sigma_1\sigma_3$	$\sigma_1\sigma_3$	$\sigma_3\sigma_1$	$-\sigma_3$	σ_3	$\sigma_3\sigma_2\sigma_1$	$\sigma_1\sigma_2\sigma_3$	σ_1	$-\sigma_1$	$\sigma_2\sigma_3$	$\sigma_3\sigma_2$	$\sigma_2\sigma_1$	$\sigma_1\sigma_2$	1	-1	σ_2	$-\sigma_2$
$\sigma_3\sigma_1$	$\sigma_3\sigma_1$	$\sigma_1\sigma_3$	σ_3	$-\sigma_3$	$\sigma_1\sigma_2\sigma_3$	$\sigma_3\sigma_2\sigma_1$	$-\sigma_1$	σ_1	$\sigma_3\sigma_2$	$\sigma_2\sigma_2$	$\sigma_1\sigma_2$	$\sigma_2\sigma_1$	-1	1	$-\sigma_2$	σ_2
$\sigma_2\sigma_3$	$\sigma_2\sigma_3$	$\sigma_3\sigma_2$	$\sigma_1\sigma_2\sigma_3$	$\sigma_3\sigma_2\sigma_1$	$-\sigma_3$	σ_3	σ_2	$-\sigma_2$	$\sigma_3\sigma_1$	$\sigma_1\sigma_3$	-1	1	$\sigma_2\sigma_1$	$\sigma_1\sigma_2$	$-\sigma_1$	σ_1
$\sigma_3\sigma_2$	$\sigma_3\sigma_2$	$\sigma_2\sigma_3$	$\sigma_3\sigma_2\sigma_1$	$\sigma_1\sigma_2\sigma_3$	σ_3	$-\sigma_3$	$-\sigma_2$	σ_2	$\sigma_1\sigma_3$	$\sigma_3\sigma_1$	1	-1	$\sigma_1\sigma_2$	$\sigma_2\sigma_1$	σ_1	$-\sigma_1$
$\sigma_1\sigma_2\sigma_3$	$\sigma_1\sigma_2\sigma_3$	$\sigma_3\sigma_2\sigma_1$	$\sigma_2\sigma_3$	$\sigma_3\sigma_2$	$\sigma_3\sigma_1$	$\sigma_1\sigma_3$	$\sigma_1\sigma_2$	$\sigma_2\sigma_1$	$-\sigma_3$	σ_3	σ_1	$-\sigma_1$	$-\sigma_2$	σ_2	-1	1
$\sigma_3\sigma_2\sigma_1$	$\sigma_3\sigma_2\sigma_1$	$\sigma_1\sigma_2\sigma_3$	$\sigma_3\sigma_2$	$\sigma_2\sigma_3$	$\sigma_1\sigma_3$	$\sigma_3\sigma_1$	$\sigma_2\sigma_1$	$\sigma_1\sigma_2$	σ_3	$-\sigma_3$	$-\sigma_1$	σ_1	σ_2	$-\sigma_2$	1	-1

The key findings from Table 1 are summarized as follows:

1. B_3 is non Abelian Group.
2. All the finite sub-groups B_3 of are:

$$\{\pm 1\}, \{1, \sigma_1\}, \{1, -\sigma_1\}, \{1, \sigma_2\}, \{1, -\sigma_2\}, \{1, \sigma_3\}, \{1, -\sigma_3\}, \{\pm 1, \pm \sigma_1\}, \{\pm 1, \pm \sigma_2\}, \{\pm 1, \pm \sigma_3\},$$

$$\{\pm 1, \pm \sigma_1\sigma_2\}, \{\pm 1, \pm \sigma_1\sigma_3\}, \{\pm 1, \pm \sigma_2\sigma_3\}, \{\pm 1, \pm \sigma_1\sigma_2\sigma_3\}, \{\pm 1, \pm \sigma_1\sigma_2, \pm \sigma_1\sigma_3, \pm \sigma_2\sigma_3\},$$

$$\{\pm 1, \pm \sigma_1, \pm \sigma_2, \pm \sigma_1\sigma_2\}, \{\pm 1, \pm \sigma_2, \pm \sigma_3, \pm \sigma_2\sigma_3\}, \{\pm 1, \pm \sigma_1, \pm \sigma_3, \pm \sigma_1\sigma_3\}, G$$

3. We define a relation \sim in B_3 such that $x \sim y$ if and only if $\exists g \in B_3$ such that $gx = y$. Clearly \sim is an equivalence relation. Hence it forms a partition on B_3 .

4. Orbit of $x = \{y \in B_3 : y \sim x\}$. Thus the Orbits are equivalence classes.

5. B_3 contains only one equivalence class.

6. The action of B_3 on itself or any other subgroup of it are:

(a) Sharply transitive as for every pair of element $x, y \in B_3 \exists$ a unique $g \in B_3$ such that $gx = y$.

(b) Faithful or Effective as different elements of B_3 induce different permutations of B_3 .

(c) Free or semi-regular as $gx = x$ for some $x \in B_3$, then $g = e = 1$.

i.e, the stabilizer of $x = Bx = \{g \in B_3 / gx = x\} = \{e = 1\}$, The kernel 'k' of the homomorphism $\phi : B_3 \rightarrow B_3$ is given by the intersection of the stabilizers Bx for all $x \in B$ which is the trivial subgroup $\{1\}$.

Note: Every free action on a non-empty set is faithful. As subgroup (subset) of B_3 is both transitive and regular, it is called a principle homogeneous space.

5. Conjugate table of the subgroup B_3

Further, we can draw conclusions on the action of B_3 on all its subgroups. We define the operation by conjugation, i.e $\Psi : B_3 \times B_3 \rightarrow B_3$ such that $\Psi(x, y) = y^x = x^{-1}yx$.

We can see that, the following are satisfied [16]:

1. $y^e = y$.
2. $(y^x)^z = z^{-1}(x^{-1}yx)z = (xz)^{-1}y(xz) = y^{xz}$.

which shows that Ψ also is a group action.

The group action of B_3 on its subgroups has some important properties which we shall discuss with the help of the conjugacy table from Table 2.

Table 2. Conjugate table

	1	-1	σ_1	$-\sigma_1$	σ_2	$-\sigma_2$	σ_3	$-\sigma_3$	$\sigma_1\sigma_2$	$\sigma_2\sigma_1$	$\sigma_1\sigma_3$	$\sigma_3\sigma_1$	$\sigma_2\sigma_3$	$\sigma_3\sigma_2$	$\sigma_1\sigma_2\sigma_3$	$\sigma_3\sigma_2\sigma_1$
1	1	-1	σ_1	$-\sigma_1$	σ_2	$-\sigma_2$	σ_3	$-\sigma_3$	$\sigma_1\sigma_2$	$\sigma_2\sigma_1$	$\sigma_1\sigma_3$	$\sigma_3\sigma_1$	$\sigma_2\sigma_3$	$\sigma_3\sigma_2$	$\sigma_1\sigma_2\sigma_3$	$\sigma_3\sigma_2\sigma_1$
-1	1	-1	σ_1	$-\sigma_1$	σ_2	$-\sigma_2$	σ_3	$-\sigma_3$	$\sigma_1\sigma_2$	$\sigma_2\sigma_1$	$\sigma_1\sigma_3$	$\sigma_3\sigma_1$	$\sigma_2\sigma_3$	$\sigma_3\sigma_2$	$\sigma_1\sigma_2\sigma_3$	$\sigma_3\sigma_2\sigma_1$
σ_1	1	-1	σ_1	$-\sigma_1$	$-\sigma_2$	σ_2	$-\sigma_3$	σ_3	$\sigma_2\sigma_1$	$\sigma_1\sigma_2$	$\sigma_3\sigma_1$	$\sigma_1\sigma_3$	$\sigma_2\sigma_3$	$\sigma_3\sigma_2$	$\sigma_1\sigma_2\sigma_3$	$\sigma_3\sigma_2\sigma_1$
$-\sigma_1$	1	-1	σ_1	$-\sigma_1$	$-\sigma_2$	σ_2	$-\sigma_3$	σ_3	$\sigma_2\sigma_1$	$\sigma_1\sigma_2$	$\sigma_3\sigma_1$	$\sigma_1\sigma_3$	$\sigma_2\sigma_3$	$\sigma_3\sigma_2$	$\sigma_1\sigma_2\sigma_3$	$\sigma_3\sigma_2\sigma_1$
σ_2	1	-1	$-\sigma_1$	σ_1	σ_2	$-\sigma_2$	$-\sigma_3$	σ_3	$\sigma_2\sigma_1$	$\sigma_1\sigma_2$	$\sigma_1\sigma_3$	$\sigma_3\sigma_1$	$\sigma_3\sigma_2$	$\sigma_2\sigma_3$	$\sigma_1\sigma_2\sigma_3$	$\sigma_3\sigma_2\sigma_1$
$-\sigma_2$	1	-1	$-\sigma_1$	σ_1	σ_2	$-\sigma_2$	$-\sigma_3$	σ_3	$\sigma_2\sigma_1$	$\sigma_1\sigma_2$	$\sigma_1\sigma_3$	$\sigma_3\sigma_1$	$\sigma_3\sigma_2$	$\sigma_2\sigma_3$	$\sigma_1\sigma_2\sigma_3$	$\sigma_3\sigma_2\sigma_1$
σ_3	1	-1	$-\sigma_1$	σ_1	$-\sigma_2$	σ_2	σ_3	$-\sigma_3$	$\sigma_1\sigma_2$	$\sigma_2\sigma_1$	$\sigma_3\sigma_1$	$\sigma_1\sigma_3$	$\sigma_3\sigma_2$	$\sigma_2\sigma_3$	$\sigma_1\sigma_2\sigma_3$	$\sigma_3\sigma_2\sigma_1$
$-\sigma_3$	1	-1	$-\sigma_1$	σ_1	$-\sigma_2$	σ_2	σ_3	$-\sigma_3$	$\sigma_1\sigma_2$	$\sigma_2\sigma_1$	$\sigma_3\sigma_1$	$\sigma_1\sigma_3$	$\sigma_3\sigma_2$	$\sigma_2\sigma_3$	$\sigma_1\sigma_2\sigma_3$	$\sigma_3\sigma_2\sigma_1$
$\sigma_1\sigma_2$	1	-1	$-\sigma_1$	σ_1	$-\sigma_2$	σ_2	σ_3	$-\sigma_3$	$\sigma_1\sigma_2$	$\sigma_2\sigma_1$	$\sigma_3\sigma_1$	$\sigma_1\sigma_3$	$\sigma_3\sigma_2$	$\sigma_2\sigma_3$	$\sigma_1\sigma_2\sigma_3$	$\sigma_3\sigma_2\sigma_1$
$\sigma_2\sigma_1$	1	-1	$-\sigma_1$	σ_1	$-\sigma_2$	σ_2	σ_3	$-\sigma_3$	$\sigma_1\sigma_2$	$\sigma_2\sigma_1$	$\sigma_3\sigma_1$	$\sigma_1\sigma_3$	$\sigma_3\sigma_2$	$\sigma_2\sigma_3$	$\sigma_1\sigma_2\sigma_3$	$\sigma_3\sigma_2\sigma_1$
$\sigma_1\sigma_3$	1	-1	$-\sigma_1$	σ_1	σ_2	$-\sigma_2$	$-\sigma_3$	σ_3	$\sigma_2\sigma_1$	$\sigma_1\sigma_2$	$\sigma_1\sigma_3$	$\sigma_3\sigma_1$	$\sigma_3\sigma_2$	$\sigma_2\sigma_3$	$\sigma_1\sigma_2\sigma_3$	$\sigma_3\sigma_2\sigma_1$
$\sigma_3\sigma_1$	1	-1	$-\sigma_1$	σ_1	σ_2	$-\sigma_2$	$-\sigma_3$	σ_3	$\sigma_2\sigma_1$	$\sigma_1\sigma_2$	$\sigma_1\sigma_3$	$\sigma_3\sigma_1$	$\sigma_3\sigma_2$	$\sigma_2\sigma_3$	$\sigma_1\sigma_2\sigma_3$	$\sigma_3\sigma_2\sigma_1$
$\sigma_2\sigma_3$	1	-1	σ_1	$-\sigma_1$	$-\sigma_2$	σ_2	$-\sigma_3$	σ_3	$\sigma_2\sigma_1$	$\sigma_1\sigma_2$	$\sigma_3\sigma_1$	$\sigma_1\sigma_3$	$\sigma_2\sigma_3$	$\sigma_3\sigma_2$	$\sigma_1\sigma_2\sigma_3$	$\sigma_3\sigma_2\sigma_1$
$\sigma_3\sigma_2$	1	-1	σ_1	$-\sigma_1$	$-\sigma_2$	σ_2	$-\sigma_3$	σ_3	$\sigma_2\sigma_1$	$\sigma_1\sigma_2$	$\sigma_3\sigma_1$	$\sigma_1\sigma_3$	$\sigma_2\sigma_3$	$\sigma_3\sigma_2$	$\sigma_1\sigma_2\sigma_3$	$\sigma_3\sigma_2\sigma_1$
$\sigma_1\sigma_2\sigma_3$	1	-1	σ_1	$-\sigma_1$	σ_2	$-\sigma_2$	σ_3	$-\sigma_3$	$\sigma_1\sigma_2$	$\sigma_2\sigma_1$	$\sigma_1\sigma_3$	$\sigma_3\sigma_1$	$\sigma_2\sigma_3$	$\sigma_3\sigma_2$	$\sigma_1\sigma_2\sigma_3$	$\sigma_3\sigma_2\sigma_1$
$\sigma_3\sigma_2\sigma_1$	1	-1	σ_1	$-\sigma_1$	σ_2	$-\sigma_2$	σ_3	$-\sigma_3$	$\sigma_1\sigma_2$	$\sigma_2\sigma_1$	$\sigma_1\sigma_3$	$\sigma_3\sigma_1$	$\sigma_2\sigma_3$	$\sigma_3\sigma_2$	$\sigma_1\sigma_2\sigma_3$	$\sigma_3\sigma_2\sigma_1$

The key findings from Table 2 are outlined as follows:

Let $B_3 = G$:

1. The Normal subgroups are,

$$\{\pm 1\}, \{\pm 1, \pm\sigma_1\}, \{\pm 1, \pm\sigma_2\}, \{\pm 1, \pm\sigma_3\},$$

$$\{\pm 1, \pm\sigma_1\sigma_2\}, \{\pm 1, \pm\sigma_1\sigma_3\}, \{\pm 1, \pm\sigma_2\sigma_3\}, \{\pm 1, \pm\sigma_1\sigma_2\sigma_3\}, \{\pm 1, \pm\sigma_1\sigma_2, \pm\sigma_1\sigma_3, \pm\sigma_2\sigma_3\},$$

$$\{\pm 1, \pm\sigma_1, \pm\sigma_2, \pm\sigma_1\sigma_2\} \{\pm 1, \pm\sigma_2, \pm\sigma_3, \pm\sigma_2\sigma_3\}, \{\pm 1, \pm\sigma_1, \pm\sigma_3, \pm\sigma_1\sigma_3\}$$

2. The Subnormal series of the above group have isomorphic refinement.

$$\{1\} \subset \{\pm 1\} \subset \{\pm 1, \pm \sigma_1\} \subset \{\pm 1, \pm \sigma_1, \pm \sigma_2, \pm \sigma_1 \sigma_2\} \subset G$$

$$\{1\} \subset \{\pm 1\} \subset \{\pm 1, \pm \sigma_2\} \subset \{\pm 1, \pm \sigma_1, \pm \sigma_2, \pm \sigma_1 \sigma_2\} \subset G$$

$$\{1\} \subset \{\pm 1\} \subset \{\pm 1, \pm \sigma_1\} \subset \{\pm 1, \pm \sigma_1, \pm \sigma_3, \pm \sigma_1 \sigma_3\} \subset G$$

$$\{1\} \subset \{\pm 1\} \subset \{\pm 1, \pm \sigma_3\} \subset \{\pm 1, \pm \sigma_1, \pm \sigma_3, \pm \sigma_1 \sigma_3\} \subset G$$

$$\{1\} \subset \{\pm 1\} \subset \{\pm 1, \pm \sigma_2\} \subset \{\pm 1, \pm \sigma_2, \pm \sigma_3, \pm \sigma_2 \sigma_3\} \subset G$$

$$\{1\} \subset \{\pm 1\} \subset \{\pm 1, \pm \sigma_3\} \subset \{\pm 1, \pm \sigma_2, \pm \sigma_3, \pm \sigma_2 \sigma_3\} \subset G$$

These are a few examples of many.

3. This shows that G is not a Simple Group as it has Normal subgroups.

4. G is Solvable.

6. Conclusion

In this paper, we have shown the total number of Normal subgroup and subnormal series of the subgroup B_3 of Clifford Algebra and by using group action and conjugation operation we given that the given subgroup B_3 is Solvable. we have taken $C(E)$, an algebra known as Clifford Algebra is constructed over a vector space E that has n -dimensions and have investigated by Group Actions the nature of its subgroups. The subgroup B_3 is a non-abelian group which has seventeen non-trivial finite subgroups with only one equivalence class and the Kernel of the homomorphism $\phi : B_3 \rightarrow B_3$ is the trivial subgroup $\{1\}$.

The Action of the subgroup B_3 with the conjugation, i.e, $\Psi : B_3 \times B_3 \rightarrow B_3$ such that $\Psi(x, y) = yx = x^{-1}yx$ has twelve Normal subgroup which implies that the subgorup B_3 is not a Simple group, further we establish that it has a subnormal series, hence we show that the subgorup B_3 is a Solvable group.

Conflict of interest

The authors declare no competing interest.

References

- [1] Hestenes D, Sobczyk G. *Clifford Algebra to Geometric Calculus: A Unified Language Formathematics and Physics*. Dordrecht: Springer; 2012.
- [2] Boi L. Clifford geometric algebras, spin manifolds, and group actions in mathematics and physics. *Advances in Applied Clifford Algebras*. 2009; 19(3): 611-656.
- [3] Delanghe R. Clifford analysis: history and perspective. *Computational Methods and Function Theory*. 2001; 1(1): 107-153.
- [4] Jayme Vaz JR, Doldao Da Rocha JR. *Clifford Analysis: History and Perspective*. New York: Oxford University Press; 2016.

- [5] Garling DJ. *Clifford Algebras: An Introduction*. Cambridge, England: Cambridge University Press; 2011.
- [6] Hestenes D. *New Foundations for Classical Mechanics*. German: Springer Science & Business Media; 2012.
- [7] Lounesto P. *Clifford Algebras and Spinors*. Cambridge, England: Cambridge University Press; 2001.
- [8] Kumar K, Goyal D, Martha S. Algebraic method for approximate solution of scattering of surface waves by thin vertical barrier over a stepped bottom topography. *Contemporary Mathematics*. 2022; 3(4): 500-513.
- [9] Riesz M. *Clifford Numbers and Spinors*. German: Springer Science & Business Media; 2013.
- [10] Shimura G. *Arithmetic and Analytic Theories of Quadratic Forms and Clifford Groups*. American: American Mathematical Soc.; 2004.
- [11] Varlamov V. Universal coverings of orthogonal groups. *Advances in Applied Clifford Algebras*. 2004; 14(1): 81-168.
- [12] Bertram W. Graded sets, graded groups, and Clifford algebras. *arXiv*. 2021. Available from: <https://doi.org/10.48550/arXiv.2109.00878>.
- [13] Hall M. *The Theory of Groups*. Mineola, New York: Courier Dover Publications; 2018.
- [14] Estaremi Y, Jabbarzadeh M. Some algebraic properties of banach space. *Contemporary Mathematics*. 2020; 1(2): 54-62. Available from: <https://doi.org/10.37256/cm.122020209>.
- [15] Khanna VK, Bhamri S. *A Course in Abstract Algebra*. India: Vikas Publishing House; 2016.
- [16] Ahmakhah N, Zarrin M. On the set of same-size conjugate classes. *Communications in Algebra*. 2019; 47(10): 3932-3938.