

Research Article

Exploring the Elzaki Transform: Unveiling Solutions to Reaction-Diffusion Equations with Generalized Composite Fractional Derivatives

Mohd Khalid¹, Ishfaq Ahmad Mallah¹, Subhash Alha¹, Ali Akgül^{2,3*}

¹Department of Mathematics, Maulana Azad National Urdu University, Gachibowli, Hyderabad-500032, India

²Department of Mathematics, Art and Science Faculty, Siirt University, 56100 Siirt, Turkey

³Department of Electronics and Communication Engineering, Saveetha School of Engineering, Chennai, Tamilnadu, India
E-mail: aliakgul00727@gmail.com

Received: 20 March 2024; **Revised:** 20 May 2024; **Accepted:** 31 May 2024

Abstract: This article investigates the use of the Elzaki transform on a generalized composite fractional derivative. To establish the framework for this inquiry, numerous essential lemmas about the Elzaki transform are presented. We successfully extract the solution to the reaction-diffusion problem using both the Elzaki and Fourier transforms, which include a generalized composite fractional derivative. We also look at special examples of the generalized equation, which helps us understand its applications and consequences better. The results show that the Elzaki transform is successful in dealing with complicated fractional differential equations, introducing new analytical approaches and solutions to the subject of fractional calculus and its applications in reaction-diffusion systems.

Keywords: elzaki transform, generalised composite fractional derivative, reaction-diffusion equation, fourier transform, fractional calculus applications

MSC: 34A08, 26A33, 35A22

1. Introduction

Fractional calculus provides a more flexible mathematical foundation and focuses on derivatives and integrals of non-integer order. Adding fractional derivatives to Reaction-diffusion equations (RDE) provides more flexibility when simulating complex systems with anomalous diffusion and memory effects. The fractional RDE has numerous applications, ranging from biology, and chemistry to physics and economics. They provide a valuable tool for modeling anomalous diffusion, subdiffusion, superdiffusion, and long-range interactions, as opposed to standard integer-order RDE [1–3]. Fractional derivatives are non-local and non-linear, making fractional RDE challenging to solve. A variety of numerical techniques, including finite difference, finite element, and spectral methods, can be used to approximate the solutions [4, 5]. In 1996, Grindrod, P. offers the conventional RDE [5].

$$\frac{\partial P}{\partial t} = D \frac{\partial^2 P}{\partial x^2} + v \cdot X(P), \quad (1)$$

where $X(P)$ is a nonlinear function representing the kinetics of a reaction and D is the diffusion coefficient.

Manne et al. [6] presented a generalization of the equation (1) as

$$\frac{\partial^2 P}{\partial t^2} + \vartheta \frac{\partial P}{\partial t} = \vartheta^2 \frac{\partial^2 P}{\partial x^2} + \lambda^2 P(x, t). \quad (2)$$

Saxena et al. [7] further generalized the RDE by illustrating with fractional derivatives

$${}_0\mathbb{D}_t^\nu P(x, t) + \vartheta \cdot {}_0\mathbb{D}_t^\mu P(x, t) = \vartheta^2 {}_{-\infty}\mathbb{D}_x^\gamma P(x, t) + \lambda^2 P(x, t) + \Theta(x, t). \quad (3)$$

which describes the diffusion transport of the quantity $P(x, t)$ in space, λ indicates the strength of the nonlinearity of the system, ϑ^2 is the diffusive constant, Θ is a constant that describes the nonlinearity in the system, $\Theta(x, t)$ is a nonlinear function for reaction kinetics. The terms ${}_0\mathbb{D}_t^\nu$ and ${}_0\mathbb{D}_t^\mu$ represent the fractional time derivatives of orders ν and μ , respectively, which reflect the memory effect and non-local features of the process. The system is nonlinear if $\nu > \mu$ and both $\Theta(x, t)$ and ϑ are nonzero. Several writers have recently researched RDEs with fractional derivatives [8–13].

Simulations of complex phenomena continue to demonstrate the utility of fractional derivatives. As a result, fractional derivatives continue to make consistent progress. This fractional derivative, as defined by Hilfer [14], incorporates the characteristics of the Riemann-Liouville fractional derivative and the Caputo fractional derivative of the same order [15]. To obtain a closed-form solution to a generalized fractional FLE equation, Garg et al. [16] defined the composition of these derivatives, allowing for different fractional orders of Riemann-Liouville and Caputo fractional derivatives. In addition, using Fourier and Sumudu transforms, Alha et al. [17] found the solution to the nonlinear RDE with a generalized composite fractional (GCF) derivative. Numerous scientific and technological disciplines employ integral transforms such as Laplace, Mellin, Fourier, Sumudu, Hankel, Elzaki, etc. [18–20]. Tarig Elzaki [21] created the Elzaki Transform as an alternative to the traditional Fourier integral for solving ordinary and partial differential equations in the time domain. This work discusses the Elzaki transform of the GCF derivative and its application to solving the reaction-diffusion equation (RDE). The uniqueness comes in extending the Elzaki transform to handle complicated GCF derivatives, which is a difficult undertaking owing to their convoluted structure. This study introduces new analytical techniques and advances the practical applications of fractional calculus in a variety of scientific domains.

2. Definition and preliminaries

Definition 1 [15] The Riemann Liouville fractional (RLF) integral of order $\nu > 0$ of a function $\Theta(t)$ is

$${}_0\mathbb{I}_t^\nu \Theta(t) = \frac{1}{\Gamma(\nu)} \int_0^t (t-u)^{\nu-1} \Theta(u) du, \quad \nu \in \mathbb{C} \text{ and } t > 0. \quad (4)$$

Definition 2 [15] The RLF derivative of order $\nu > 0$ of a function $\Theta(t)$ is

$${}_0\mathbb{D}_t^\nu \Theta(t) = \frac{1}{\Gamma(\kappa - \nu)} \frac{d^\kappa}{dt^\kappa} \int_0^t (t-u)^{\kappa-\nu-1} \Theta(u) du, \quad \kappa - 1 < \nu < \kappa, \quad \kappa \in \mathbb{N}. \quad (5)$$

Definition 3 [15] Caputo fractional (CF) derivative of order $\nu > 0$ of a function $\Theta(t)$ is

$${}_0^C \mathbb{D}_t^\nu \Theta(t) = \frac{1}{\Gamma(\kappa - \nu)} \int_0^t (t-u)^{\kappa-\nu-1} \Theta^{(\kappa)}(u) du, \quad \kappa - 1 < \nu < \kappa, \quad \kappa \in \mathbb{N}. \quad (6)$$

Definition 4 [7, 22] Weyl fractional (WF) derivative of order $\nu > 0$ is defined as

$${}_{-\infty} \mathbb{D}_t^\nu \Theta(t) = \frac{1}{\Gamma(\kappa - \nu)} \frac{d^\kappa}{dt^\kappa} \int_{-\infty}^t (t-u)^{\kappa-\nu-1} \Theta(u) du, \quad \kappa - 1 < \nu < \kappa, \quad \kappa \in \mathbb{N}. \quad (7)$$

The modified Fourier integral transform (FIT) of the operator (7) given by Metzler and Klafter [23] is

$$F \{ {}_{-\infty} \mathbb{D}_t^\nu \Theta(t) \} = -\kappa^\nu \Theta^*(\kappa), \quad (8)$$

where the FIT is defined by means of the integral equation

$$\Theta^*(\kappa) = \int_{-\infty}^{\infty} \Theta(t) \exp(i\kappa t) dt. \quad (9)$$

Definition 5 [14] Hilfer fractional (HF) derivative of order ν and μ of a function $\Theta(t)$ is defined as

$${}_0 \mathbb{D}_t^{\nu, \mu} \Theta(t) = \left({}_0 \mathbb{I}_t^{\mu(1-\nu)} \frac{d}{dt} ({}_0 \mathbb{I}_t^{(1-\nu)(1-\mu)} \Theta(t)) \right), \quad (10)$$

where $0 < \nu \leq 1$, and $0 \leq \mu \leq 1$.

Definition 6 [16] The GCF derivative of a function $\Theta(t)$ is defined as

$${}_0 \mathbb{D}_t^{\nu, \mu; \alpha} \Theta(t) = \left({}_0 \mathbb{I}_t^{\alpha(\kappa-\mu)} \frac{d^\kappa}{dt^\kappa} ({}_0 \mathbb{I}_t^{(1-\alpha)(\kappa-\nu)} \Theta(t)) \right), \quad (11)$$

where $\kappa - 1 < \nu$, $\mu \leq \kappa$; $0 \leq \alpha \leq 1$ and $\kappa \in \mathbb{N}$.

For $\alpha = 0$ and $\alpha = 1$, the GCF derivative (11) simplifies to the RLF derivative of order ν (5) and the CF derivative of order μ (6). For $\nu = \mu$, the GCF derivative is same as the ν -order, α -type derivative of HF derivative (10).

Definition 7 [24] A generalization of the Mittag-Leffler function proposed by Prabhakar is

$$E_{\nu, \mu}^\gamma(t) = \sum_{j=0}^{\infty} \frac{(\gamma)_j}{\Gamma(\nu j + \mu)} \cdot \frac{t^j}{j!}, \quad (12)$$

where ν , μ and γ are the complex numbers with $Re(\nu)$, $Re(\mu) > 0$.

Definition 8 [21] The Elzaki integral transform (EIT) denoted by $T(s)$ for the function $\Theta(t)$ is expressed as follows:

$$E[\Theta(t), s] = s^2 \int_0^\infty \exp(-t) \Theta(st) dt = s \int_0^\infty \exp(-\frac{t}{s}) \Theta(t) dt, \quad s \in (\ell_1, \ell_2),$$

for the set of functions

$$\mathcal{A} = \left\{ \Theta(t) \mid \exists M, \ell_1, \ell_2 > 0, |\Theta(t)| < M \exp\left(\frac{|t|}{\ell_j}\right), \text{ if } t \in (-1)^j \times [0, \infty) \right\}.$$

In the upcoming sections, it will help to know about some of the EIT's most important features

Proposition 1 [25] If $M(s)$ and $N(s)$ are the EIT's for $\Theta(t)$ and $\varphi(t)$ respectively, then their convolution is defined as

$$E[\Theta(t) * \varphi(t), s] = \frac{1}{s} M(s)N(s), \quad (13)$$

or equivalently

$$E^{-1}\left[\frac{1}{s} M(s)N(s), t\right] = (\Theta(t) * \varphi(t)), \quad (14)$$

where

$$(\Theta(t) * \varphi(t)) = \frac{1}{s} \int_0^t \Theta(u)\varphi(t-u)du.$$

Lemma 1 [26] The EIT of RLF integral of order ν is defined by

$$E[{}_0\mathbb{I}_t^\nu \Theta(t), s] = s^\nu T(s), \quad \Re(\nu) > 0. \quad (15)$$

Lemma 2 [27] The EIT of m^{th} derivative $\Theta^{(m)}(t)$ is defined by

$$E[\Theta^{(m)}(t), s] = s^{-m} T(s) - \sum_{j=0}^{m-1} s^{j-m+2} \Theta^{(j)}(0), \quad m \geq 0. \quad (16)$$

Lemma 3 [28] The EIT of the RLF and CF derivatives of order ν are defined as

$$E[{}_0\mathbb{D}_t^\nu \Theta(t), s] = s^{-\nu} T(s) - \sum_{j=0}^{\kappa-1} s^{-(j-2)} [\mathbb{D}^{\nu-j} \Theta(t)]_{t=0}$$

and

$$E[{}_0^C\mathbb{D}_t^\nu \Theta(t), s] = s^{-\nu} T(s) - \sum_{j=0}^{\kappa-1} s^{2-\nu+j} [\mathbb{D}^\kappa \Theta(t)]_{t=0},$$

where $\kappa - 1 \leq \nu < \kappa$, $\kappa \in \mathbb{N}$.

Lemma 4 [29] In the complex plane \mathbb{C} , the following equality holds for the inverse of EIT:

$$E^{-1} \left[u^{\gamma+1} (1 - \rho u^\mu)^{-\delta} \right] = t^{\gamma-1} E_{\mu, \gamma}^\delta (\rho u^\mu), \quad (17)$$

where μ, γ, δ and $\rho \in \mathbb{C}$ with $Re(\mu) > 0$, $Re(\gamma) > 0$.

3. Main results

Lemma 5 Let $0 \leq \alpha \leq 1$, $\nu - \alpha(\nu - \mu) > \omega - \alpha(\omega - \delta)$ and $\kappa \in \mathbb{N}$, therefore, the succeeding equivalence holds true

$$\begin{aligned} & E^{-1} \left[\frac{1}{s^{-1}(s^{\alpha(\nu-\mu)-\nu} + \vartheta \cdot s^{\alpha(\omega-\delta)-\omega} + \eta)} \right] \\ &= \sum_{r=0}^{\infty} (-\eta)^r t^{\{v-\alpha(v-\mu)\}(r+1)-1} \times E_{\nu-\alpha(v-\mu)-\omega+\alpha(\omega-\delta), \{v-\alpha(v-\mu)\}(r+1)}^{(r+1)} \left(-\vartheta \cdot t^{v-\alpha(v-\mu)-\omega+(\omega-\delta)} \right), \end{aligned} \quad (18)$$

where $\nu, \mu > \kappa - 1$; $\kappa > \omega, \delta$.

Proof. For LHS of (18), we consider

$$\begin{aligned} & \frac{1}{s^{-1}(s^{\alpha(\nu-\mu)-\nu} + \vartheta \cdot s^{\alpha(\omega-\delta)-\omega} + \eta)} \\ &= \left[\frac{1}{s^{-1}(s^{\alpha(\nu-\mu)-\nu} + \vartheta \cdot s^{\alpha(\omega-\delta)-\omega})} \right] \times \left[1 + \frac{\eta}{s^{\alpha(\nu-\mu)-\nu} + \vartheta \cdot s^{\alpha(\omega-\delta)-\omega}} \right]^{-1} \\ &= \left[\frac{s}{s^{\alpha(\nu-\mu)-\nu} + \vartheta \cdot s^{\alpha(\omega-\delta)-\omega}} \right] \times \sum_{r=0}^{\infty} \frac{(-\eta)^r}{[s^{\alpha(\nu-\mu)-\nu} + \vartheta \cdot s^{\alpha(\omega-\delta)-\omega}]^r} \\ &= \sum_{r=0}^{\infty} (-\eta)^r \cdot s^{\{v-\alpha(v-\mu)\}(r+1)+1} \times \left[1 + \vartheta \cdot s^{v-\alpha(v-\mu)-\omega+\alpha(\omega-\delta)} \right]^{-(r+1)}. \end{aligned} \quad (19)$$

Now, taking the inverse of EIT on (19) and using (17), we arrive at (18). □

Lemma 6 Let $0 \leq \alpha \leq 1$, $\nu - \alpha(\nu - \mu) > \omega - \alpha(\omega - \delta)$ and $\kappa \in \mathbb{N}$, therefore, the succeeding equivalence holds true

$$\begin{aligned}
& E^{-1} \left[\frac{s^{\alpha(\kappa-\mu)-\kappa}}{(s^{\alpha(v-\mu)-v} + \vartheta \cdot s^{\alpha(\omega-\delta)-\omega} + \eta)} \right] \\
&= \sum_{r=0}^{\infty} (-\eta)^r t^{\alpha(\kappa+r\mu)+v(r+1)(1-\alpha)-\kappa-2} \times E_{v-\alpha(v-\mu)-\omega+\alpha(\omega-\delta), \alpha(\kappa+r\mu)+v(r+1)(1-\alpha)-\kappa-1}^{(r+1)} \\
&\quad \left(-\vartheta \cdot t^{v-\alpha(v-\mu)-\omega+\alpha(\omega-\delta)} \right),
\end{aligned} \tag{20}$$

where $v, \mu > \kappa - 1$; $\kappa > \omega, \delta$.

Proof. To illustrate (20), we consider

$$\begin{aligned}
\frac{s^{\alpha(\kappa-\mu)-\kappa}}{(s^{\alpha(v-\mu)-v} + \vartheta \cdot s^{\alpha(\omega-\delta)-\omega} + \eta)} &= \frac{s^{\alpha(\kappa-\mu)-\kappa}}{s^{\alpha(v-\mu)-v} + \vartheta \cdot s^{\alpha(\omega-\delta)-\omega}} \left(1 + \frac{\eta}{s^{\alpha(v-\mu)-v} + \vartheta \cdot s^{\alpha(\omega-\delta)-\omega}} \right)^{-1} \\
&= \sum_{r=0}^{\infty} (-\eta)^r s^{\alpha(\kappa-\mu)-\kappa-\{\alpha(v-\mu)-v\}(r+1)} \left(1 + \vartheta \cdot s^{v-\alpha(v-\mu)-\omega+\alpha(\omega-\delta)} \right)^{-(r+1)}.
\end{aligned} \tag{21}$$

Now, taking the inverse of EIT on (21) and using (17), we arrive at (20). □

Similarly, we can write the following lemma:

Lemma 7 Let $0 \leq \alpha \leq 1$, $v - \alpha(v - \mu) > \omega - \alpha(\omega - \delta)$ and $\kappa \in \mathbb{N}$, therefore, the succeeding equivalence holds true.

$$\begin{aligned}
& E^{-1} \left[\frac{s^{\alpha(\kappa-\delta)-\kappa}}{(s^{\alpha(v-\mu)-v} + \vartheta \cdot s^{\alpha(\omega-\delta)-\omega} + \eta)} \right] \\
&= \sum_{r=0}^{\infty} (-\eta)^r t^{\kappa(\alpha-1)+(r+1)\{v(1-\alpha)+\alpha\mu\}-\alpha\delta-2} \\
&\quad \times E_{v-\alpha(v-\mu)-\omega+\alpha(\omega-\delta), \kappa(\alpha-1)+(r+1)\{v(1-\alpha)+\alpha\mu\}-\alpha\delta-1}^{(r+1)} \left(-\vartheta \cdot t^{v-\alpha(v-\mu)-\omega+\alpha(\omega-\delta)} \right),
\end{aligned} \tag{22}$$

where $v, \mu > \kappa - 1$; $\kappa > \omega, \delta$.

3.1 EIT of GCF derivative

Theorem 8 Let the EIT of the function $\Theta(t)$ denoted by $T(s)$, then the EIT of the GCF derivative ${}_0\mathbb{D}_t^{v, \mu; \alpha} \Theta(t)$ is defined by

$$E[{}_0\mathbb{D}_t^{v, \mu; \alpha} \Theta(t), s] = s^{\alpha(v-\mu)-v} T(s) - \sum_{j=0}^{\kappa-1} s^{\alpha(\kappa-\mu)-\kappa+j+2} \left[\left(\mathbb{D}^j \left({}_0\mathbb{I}_t^{(1-\alpha)(\kappa-v)} \right) \Theta(t) \right) \Big|_{t=0} \right], \tag{23}$$

where $\kappa - 1 < \nu$, $\mu \leq \kappa$ with $\kappa \in \mathbb{N}$.

Proof. Suppose $\varphi(t) = \mathbb{D}^\kappa {}_0\mathbb{I}_t^{(1-\alpha)(\kappa-\nu)} \Theta(t) = \mathbb{D}^\kappa \varphi(t)$ for (11) and using (15), we get

$$E [{}_0\mathbb{D}_t^{\nu, \mu; \alpha} \Theta(t), s] = E [{}_0\mathbb{I}_t^{\alpha(\kappa-\mu)} \varphi(t), s] = s^{\alpha(\kappa-\mu)} E[\varphi(t), s] = s^{\alpha(\kappa-\mu)} E[\mathbb{D}^\kappa \varphi(t), s], \quad (24)$$

where $\varphi(t) = {}_0\mathbb{I}_t^{(1-\alpha)(\kappa-\nu)} \Theta(t)$, using (16) in (24), we get

$$E [{}_0\mathbb{D}_t^{\nu, \mu; \alpha} \Theta(t), s] = s^{\alpha(\kappa-\mu)-\kappa} E [{}_0\mathbb{I}_t^{(1-\alpha)(\kappa-\nu)} \Theta(t), s] - \sum_{j=0}^{\kappa-1} s^{\alpha(\kappa-\mu)-\kappa+j+2} \left[\left(\mathbb{D}^j \left({}_0\mathbb{I}_t^{(1-\alpha)(\kappa-\nu)} \Theta \right) (t) \right) \Big|_{t=0} \right], \quad (25)$$

again using (15) in (25), we get

$$E [{}_0\mathbb{D}_t^{\nu, \mu; \alpha} \Theta(t), s] = s^{\alpha(\nu-\mu)-\nu} \times s^{(1-\alpha)(\kappa-\nu)} E[\Theta(t), s] - \sum_{j=0}^{\kappa-1} s^{\alpha(\kappa-\mu)-\kappa+j+2} \left[\left(\mathbb{D}^j \left({}_0\mathbb{I}_t^{(1-\alpha)(\kappa-\nu)} \Theta \right) (t) \right) \Big|_{t=0} \right], \quad (26)$$

we arrive at (23). □

3.2 The GCF derivative and its solution to the RDE

Theorem 9 For $0 \leq \alpha \leq 1$; $\kappa - 1 < \nu$, μ ; ω , $\delta \leq \kappa$, $\kappa \in \mathbb{N}$ such that $\nu > \omega$, $\delta < \mu$. Consider the fractional RDE

$${}_0\mathbb{D}_t^{\nu, \mu; \alpha} P(x, t) + \vartheta \cdot {}_0\mathbb{D}_t^{\omega, \delta; \alpha} P(x, t) = \vartheta^2 {}_{-\infty}\mathbb{D}_x^\gamma P(x, t) + \lambda^2 P(x, t) + \Theta(x, t), \quad (27)$$

with the initial conditions

$$\begin{cases} \mathbb{D}^j {}_0\mathbb{I}_t^{(1-\alpha)(\kappa-\nu)} P(x, t) \Big|_{t=0} = \varphi_1(x) \\ \mathbb{D}^j {}_0\mathbb{I}_t^{(1-\alpha)(\kappa-\omega)} P(x, t) \Big|_{t=0} = \varphi_2(x) \end{cases} ; \quad j = 0, 1, \dots, \kappa - 1, \quad (28)$$

where ϑ is a diffusion coefficient, α is a constant that represents the non-linearity of the system, and Θ is a non-linear function for reaction kinetics than the solution of (27) corresponding to $P(x, t)$ is as follows

$$\begin{aligned}
P(x, t) &= \sum_{j=0}^{\kappa-1} \sum_{r=0}^{\infty} \frac{(-\eta)^r}{\sqrt{2\pi}} \int_{-\infty}^{\infty} t^{\alpha(\kappa+r\mu)+v(r+1)(1-\alpha)-\kappa+j} \varphi_1^*(N) \exp(-iNx) \\
&\quad \times E_{v-\alpha(v-\mu)-\omega+\alpha(\omega-\delta), \alpha(\kappa+r\mu)+v(r+1)(1-\alpha)-\kappa+j+1}^{r+1} \left(-\vartheta \cdot t^{v-\alpha(v-\mu)-\omega+\alpha(\omega-\delta)} \right) dN \\
&\quad + \sum_{j=0}^{\kappa-1} \sum_{r=0}^{\infty} \vartheta \frac{(-\eta)^r}{\sqrt{2\pi}} \int_{-\infty}^{\infty} t^{\kappa(\alpha-1)+(r+1)(v-\alpha(v-\mu))-\alpha\delta+j} \varphi_2^*(N) \exp(-iNx) \\
&\quad \times E_{v-\alpha(v-\mu)-\omega+\alpha(\omega-\delta), \kappa(\alpha-1)+(r+1)(v-\alpha(v-\mu))-\alpha\delta+j+1}^{r+1} \left(-\vartheta \cdot t^{v-\alpha(v-\mu)-\omega+\alpha(\omega-\delta)} \right) dN \\
&\quad + \sum_{r=0}^{\infty} \frac{(-\eta)^r}{\sqrt{2\pi}} \int_0^t u^{\{v-\alpha(v-\mu)\}(r+1)-1} \int_{-\infty}^{\infty} \Theta^*(N, t-u) \exp(-iNx) \\
&\quad \times E_{v-\alpha(v-\mu)-\omega+\alpha(\omega-\delta), \{v-\alpha(v-\mu)\}(r+1)}^{r+1} \left(-\vartheta \cdot t^{v-\alpha(v-\mu)-\omega+\alpha(\omega-\delta)} \right) dudN.
\end{aligned} \tag{29}$$

Proof. If we use the EIT on both sides of (27) with respect to t , we get

$$\begin{aligned}
&E[{}_0\mathbb{D}_t^{v, \mu; \alpha} P(x, t), s] + E[\vartheta \cdot {}_0\mathbb{D}_t^{\omega, \delta; \alpha} P(x, t), s] \\
&= \vartheta^2 E[-\infty\mathbb{D}_x^{\gamma} P(x, t), s] + \lambda^2 E[P(x, t), s] + E[\Theta(x, t)],
\end{aligned} \tag{30}$$

using (23) and (28) in (30), we have

$$\begin{aligned}
&s^{\alpha(v-\mu)-v} \bar{P}(x, s) - \sum_{j=0}^{\kappa-1} s^{\alpha(\kappa-\mu)-\kappa+j+2} \left[\mathbb{D}^j \left({}_0\mathbb{I}_t^{(1-\alpha)(\kappa-v)} P(x, t) \right) \right] \Big|_{t=0} \\
&\quad + \vartheta \cdot s^{\alpha(\omega-\delta)-\omega} \bar{P}(x, s) - \vartheta \cdot \sum_{j=0}^{\kappa-1} s^{\alpha(\kappa-\delta)-\kappa+j+2} \left[\mathbb{D}^j \left({}_0\mathbb{I}_t^{(1-\alpha)(\kappa-\omega)} P(x, t) \right) \right] \Big|_{t=0} \\
&= \vartheta^2 {}_{-\infty}\mathbb{D}_x^{\gamma} \bar{P}(x, s) + \lambda^2 \bar{P}(x, s) + \bar{\Theta}(x, s) \\
&\quad s^{\alpha(v-\mu)-v} \bar{P}(x, s) - \sum_{j=0}^{\kappa-1} s^{\alpha(\kappa-\mu)-\kappa+j+2} \varphi_1(x) + \vartheta \cdot s^{\alpha(\omega-\delta)-\omega} \bar{P}(x, s) \\
&\quad - \vartheta \cdot \sum_{j=0}^{\kappa-1} s^{\alpha(\kappa-\delta)-\kappa+j+2} \varphi_2(x)
\end{aligned}$$

$$= \vartheta^2 \mathbb{D}_x^\gamma \bar{P}(x, s) + \lambda^2 \bar{P}(x, s) + \bar{\Theta}(x, s). \quad (31)$$

Now, by applying FIT to both ends of (31) with respect to x and employing (7), we obtain

$$\begin{aligned} s^{\alpha(v-\mu)-v} \bar{P}^*(N, s) - \sum_{j=0}^{\kappa-1} s^{\alpha(\kappa-\mu)-\kappa+j+2} \varphi_1^*(N) + \vartheta \cdot s^{\alpha(\omega-\delta)-\omega} \bar{P}^*(N, s) - \vartheta \cdot \sum_{j=0}^{\kappa-1} s^{\alpha(\kappa-\delta)-\kappa+j+2} \varphi_2^*(N) \\ = -\vartheta^2 |\kappa|^\gamma \bar{P}^*(N, s) + \lambda^2 \bar{P}^*(N, s) + \bar{\Theta}^*(N, s), \end{aligned} \quad (32)$$

to solve $\bar{P}^*(N, s)$, above corresponds to

$$\begin{aligned} \bar{P}^*(N, s) = \sum_{j=0}^{\kappa-1} \varphi_1^*(N) \frac{s^{\alpha(\kappa-\mu)-\kappa+j+2}}{s^{\alpha(v-\mu)-v} + \vartheta \cdot s^{\alpha(\omega-\delta)-\omega} + \eta} + \sum_{j=0}^{\kappa-1} \vartheta \cdot \varphi_2^*(N) \frac{s^{\alpha(\kappa-\delta)-\kappa+j+2}}{s^{\alpha(v-\mu)-v} + \vartheta \cdot s^{\alpha(\omega-\delta)-\omega} + \eta} \\ + \frac{\bar{\Theta}^*(N, s)}{s^{\alpha(v-\mu)-v} + \vartheta \cdot s^{\alpha(\omega-\delta)-\omega} + \eta}, \end{aligned} \quad (33)$$

where $\eta = \vartheta^2 |\kappa|^\gamma - \lambda^2$. Now, using (18), (22), and the EIT convolution (14), we obtained by applying the inverse EIT on both ends of (33)

$$\begin{aligned} P^*(N, t) = \sum_{j=0}^{\kappa-1} \varphi_1^*(N) \sum_{r=0}^{\infty} (-\eta)^r t^{\alpha(\kappa+r\mu)+v(r+1)(1-\alpha)-\kappa+j} \\ \times E_{v-\alpha(v-\mu)-\omega+\alpha(\omega-\delta), \alpha(\kappa+r\mu)+v(r+1)(1-\alpha)-\kappa+j+1}^{r+1} \left(-\vartheta \cdot t^{v-\alpha(v-\mu)-\omega+\alpha(\omega-\delta)} \right) \\ + \sum_{j=0}^{\kappa-1} \vartheta \cdot \varphi_2^*(N) \sum_{r=0}^{\infty} (-\eta)^r t^{\kappa(\alpha-1)+(r+1)\{v(1-\alpha)+\alpha\mu\}-\alpha\delta+j} \\ \times E_{v-\alpha(v-\mu)-\omega+\alpha(\omega-\delta), \kappa(\alpha-1)+(r+1)\{v(1-\alpha)+\alpha\mu\}-\alpha\delta+j+1}^{r+1} \left(-\vartheta \cdot t^{v-\alpha(v-\mu)-\omega+\alpha(\omega-\delta)} \right) \\ + \sum_{r=0}^{\infty} (-\eta)^r \int_0^t \Theta^*(N, t-u) u^{\{v-\alpha(v-\mu)\}(r+1)-1} \\ \times E_{v-\alpha(v-\mu)-\omega+\alpha(\omega-\delta), \{v-\alpha(v-\mu)\}(r+1)}^{r+1} \left(-\vartheta \cdot t^{v-\alpha(v-\mu)-\omega+\alpha(\omega-\delta)} \right) du, \end{aligned} \quad (34)$$

when we do an inverse FIT on both ends of (34), we get (29). □

3.3 Special cases

Corollary 1 For $\alpha = 0$, the GCF-RDE (27) reduces into the RDE with RLF's derivative, that is

$${}_0\mathbb{D}_t^\nu P(x, t) + \vartheta \cdot {}_0\mathbb{D}_t^\omega P(x, t) = \vartheta^2 {}_{-\infty}\mathbb{D}_x^\gamma P(x, t) + \lambda^2 P(x, t) + \Theta(x, t), \quad (35)$$

where $\kappa - 1 < \nu$; $\omega \leq \kappa$; $\nu > \omega$; $\kappa \in \mathbb{N}$; ${}_0\mathbb{D}_t^\nu$ and ${}_0\mathbb{D}_t^\omega$ are fractional derivatives in RLF's sense with the initial conditions

$$\begin{cases} \mathbb{D}_t^j {}_0\mathbb{I}_t^{(1-\alpha)(\kappa-\nu)} P(x, t)|_{t=0} = \varphi_1(x) \\ \mathbb{D}_t^j {}_0\mathbb{I}_t^{(1-\alpha)(\kappa-\omega)} P(x, t)|_{t=0} = \varphi_2(x) \end{cases}; \quad j = 0, 1, \dots, \kappa - 1. \quad (36)$$

Given initial conditions (36), the solution of (35) is given by

$$\begin{aligned} P(x, t) = & \sum_{j=0}^{\kappa-1} \sum_{r=0}^{\infty} \frac{(-\eta)^r}{\sqrt{2\pi}} \int_{-\infty}^{\infty} t^{\nu(r+1)-\kappa+j} \varphi_1^*(N) \exp(-iNx) \\ & \times E_{\nu-\omega, \nu(r+1)-\kappa+j+1}^{r+1} (-\vartheta \cdot t^{\nu-\omega}) dN \\ & + \sum_{j=0}^{\kappa-1} \sum_{r=0}^{\infty} \vartheta \frac{(-\eta)^r}{\sqrt{2\pi}} \int_{-\infty}^{\infty} t^{\nu(r+1)-\kappa+j} \varphi_2^*(N) \exp(-iNx) \\ & \times E_{\nu-\omega, \nu(r+1)-\kappa+j+1}^{r+1} (-\vartheta \cdot t^{\nu-\omega}) dN \\ & + \sum_{r=0}^{\infty} \frac{(-\eta)^r}{\sqrt{2\pi}} \int_0^t u^{\{\nu(r+1)-1\}} \int_{-\infty}^{\infty} \Theta^*(N, t-u) \exp(-iNx) \\ & \times E_{\nu-\omega, \nu(r+1)}^{r+1} (-\vartheta \cdot t^{\nu-\alpha(\nu-\mu)-\omega+\alpha(\omega-\delta)}) dudN. \end{aligned} \quad (37)$$

Corollary 2 For $\alpha = 1$, the GCF-RDE (27) reduces into a RDE with CF's derivative, that is

$${}_0\mathbb{D}_t^\mu P(x, t) + \vartheta \cdot {}_0\mathbb{D}_t^\delta P(x, t) = \vartheta^2 {}_{-\infty}\mathbb{D}_x^\gamma P(x, t) + \lambda^2 P(x, t) + \Theta(x, t), \quad (38)$$

where $\kappa - 1 < \mu$, $\delta \leq \kappa$; $\mu > \delta$, $\kappa \in \mathbb{R}$; ${}_0\mathbb{D}_t^\mu$ and ${}_0\mathbb{D}_t^\delta$ are fractional derivatives in CF sense with the initial conditions

$$\mathbb{D}_0^j P(x, t)|_{t=0} = \varphi(x), \quad j = 0, 1, \dots, \kappa - 1. \quad (39)$$

The expression provides the solution of (38) with initial conditions (39)

$$\begin{aligned}
P(x, t) = & \sum_{j=0}^{\kappa-1} \sum_{r=0}^{\infty} \frac{(-\eta)^r}{\sqrt{2\pi}} \int_{-\infty}^{\infty} t^{\mu r+j} \varphi_1^*(N) \exp(-iNx) \times E_{\mu-\delta, \mu r+j+1}^{r+1}(-\vartheta \cdot t^{\mu-\delta}) dN \\
& + \sum_{j=0}^{\kappa-1} \sum_{r=0}^{\infty} \vartheta \frac{(-\eta)^r}{\sqrt{2\pi}} \int_{-\infty}^{\infty} t^{\mu(r+1)-\delta+j} \varphi_2^*(N) \exp(-iNx) \\
& \times E_{\mu-\delta, \mu(r+1)-\delta+j+1}^{r+1}(-\vartheta \cdot t^{\mu-\delta}) dN \\
& + \sum_{r=0}^{\infty} \frac{(-\eta)^r}{\sqrt{2\pi}} \int_0^t u^{\mu(r+1)-1} \int_{-\infty}^{\infty} \Theta^*(N, t-u) \exp(-iNx) \times E_{\mu-\delta, \mu(r+1)}^{r+1}(-\vartheta \cdot t^{\mu-\delta}) dudN.
\end{aligned} \tag{40}$$

In particular, if we take $\kappa = 1$ in (38)-(40), we get a form of fractional RDE studied by Saxena et al. [7] and Gupta and Sharma [30].

Corollary 3 For $\nu = \mu$; $\omega = \delta$, the GCF-RDE (27) reduces into a RDE with HF's derivative, that is

$${}_0\mathbb{D}_t^{\nu, \alpha} P(x, t) + \vartheta \cdot {}_0\mathbb{D}_t^{\omega, \alpha} P(x, t) = \vartheta_{-\infty}^2 \mathbb{D}_x^{\gamma} P(x, t) + \lambda^2 P(x, t) + \Theta(x, t), \tag{41}$$

with the initial conditions

$$\begin{cases} \mathbb{D}_0^j {}_0\mathcal{I}_t^{(1-\alpha)(\kappa-\nu)} P(x, t)|_{t=0} = \varphi_1(x) \\ \mathbb{D}_0^j {}_0\mathcal{I}_t^{(1-\alpha)(\kappa-\omega)} P(x, t)|_{t=0} = \varphi_2(x) \end{cases} ; j = 0, 1, \dots, \kappa-1, \tag{42}$$

where $\kappa-1 < \nu$, $\omega \leq \kappa$ and $\kappa \in \mathbb{N}$ such that $0 \leq \alpha \leq 1$. The answer that corresponds to this problem can be found by

$$\begin{aligned}
P(x, t) = & \sum_{j=0}^{\kappa-1} \sum_{r=0}^{\infty} \frac{(-\eta)^r}{\sqrt{2\pi}} \int_{-\infty}^{\infty} t^{\nu(r+1-\alpha)+\kappa(\alpha-1)+j} \varphi_1^*(N) \exp(-iNx) \\
& \times E_{\nu-\omega, \nu(r+1-\alpha)+\kappa(\alpha-1)+j+1}^{r+1}(-\vartheta \cdot t^{\nu-\omega}) dN \\
& + \sum_{j=0}^{\kappa-1} \sum_{r=0}^{\infty} \vartheta \frac{(-\eta)^r}{\sqrt{2\pi}} \int_{-\infty}^{\infty} t^{\kappa(\alpha-1)+\nu(r+1)-\alpha\omega+j} \varphi_2^*(N) \exp(-iNx) \\
& \times E_{\nu-\omega, \kappa(\alpha-1)+\nu(r+1)-\alpha\omega+j+1}^{r+1}(-\vartheta \cdot t^{\nu-\omega}) dN \\
& + \sum_{r=0}^{\infty} \frac{(-\eta)^r}{\sqrt{2\pi}} \int_0^t u^{\nu(r+1)-1} \int_{-\infty}^{\infty} \Theta^*(N, t-u) \exp(-iNx)
\end{aligned} \tag{43}$$

$$\times E_{\nu-\omega, \nu(r+1)}^{r+1} (-\vartheta \cdot t^{\nu-\omega}) dudN$$

Specifically, the Alkahtani et al. [8] studied FDE obtained in this scenario for $\kappa = 1$.

4. Conclusion

By deriving the inverse Elzaki transform and introducing multiple lemmas related to the generalized composite fractional derivative, this study has made significant contributions to the discipline. Researchers have effectively solved a nonlinear reaction-diffusion equation incorporating the generalized composite fractional derivative by using the Elzaki and Fourier transforms. In addition, some special cases of the general equation have been discussed. Furthermore, the provided approach is potentially applicable to different forms of fractional derivatives, implying larger value in a variety of scenarios. Future research might include extending the Elzaki transform to additional fractional derivatives and complicated systems, creating numerical methods to supplement analytical answers, and investigating multidisciplinary applications in physics, biology, and engineering.

Acknowledgments

The authors are very thankful to the editor and anonymous referees for their critical comments and suggestions to improve this paper to this current form.

Conflict of interest

The authors declare no competing financial interest.

References

- [1] Chen J, Liu F, Anh V. Analytical solution for the time-fractional telegraph equation by the method of separating variables. *Journal of Mathematical Analysis and Applications*. 2008; 338(2): 1364-1377.
- [2] Gorenflo R, Mainardi F, Moretti D, Paradisi P. Time fractional diffusion: a discrete random walk approach. *Nonlinear Dynamics*. 2002; 29: 129-143.
- [3] Khalid M, Alha S. New generalized integral transform on Hilfer-Prabhakar fractional derivatives and its applications. *International Journal of Dynamics and Control*. 2024; 12(1): 24-31.
- [4] Bessonov N, Bocharov G, Meyerhans A, Popov V, Volpert V. Nonlocal reaction-diffusion model of viral evolution: Emergence of virus strains. *Mathematics*. 2020; 8(1): 117.
- [5] Grindrod P. *The Theory and Applications of Reaction-Diffusion Equations: Patterns and Waves*. New York: Oxford University Press; 1996.
- [6] Manne KK, Hurd AJ, Kenkre VM. Nonlinear waves in reaction-diffusion systems: The effect of transport memory. *Physical Review E*. 2000; 61(4): 4177.
- [7] Saxena RK, Mathai AM, Haubold HJ. Reaction-diffusion systems and nonlinear waves. *Astrophysics and Space Science*. 2006; 305: 297-303.
- [8] Alkahtani B, Gulati V, Kılıçman A. Application of Sumudu transform in generalized fractional reaction-diffusion equation. *International Journal of Applied and Computational Mathematics*. 2016; 2: 387-394.
- [9] Fazli H, Bahrami F. On the steady solutions of fractional reaction-diffusion equations. *Filomat*. 2017; 31(6): 1655-1664.

- [10] Viana A. A local theory for a fractional reaction-diffusion equation. *Communications in Contemporary Mathematics*. 2019; 21(6): 1850033.
- [11] Yao SW, Arqub OA, Tayebi S, Osman MS, Mahmoud W, Inc M, et al. A novel collective algorithm using cubic uniform spline and finite difference approaches to solving fractional diffusion singular wave model through damping-reaction forces. *Fractals*. 2023; 31(4): 2340069.
- [12] Chu Y, Rashid S, Kubra KT, Inc M, Hammouch Z, Osman MS. Analysis and numerical computations of the multi-dimensional, time-fractional model of navier-stokes equation with a new integral transformation. *CMES-Computer Modeling in Engineering & Sciences*. 2023; 136(3): 3025.
- [13] Shi L, Rashid S, Sultana S, Khalid A, Agarwal P, Osman MS. Semi-analytical view of time-fractional PDES with proportional delays pertaining to index and Mittag-Leffler memory interacting with hybrid transforms. *Fractals*. 2023; 31(4): 2340071.
- [14] Hilfer R. *Applications of Fractional Calculus in Physics*. World scientific; 2000.
- [15] Kilbas AA, Srivastava HM, Trujillo JJ. *Theory and Applications of Fractional Differential Equations*. Elsevier; 2006.
- [16] Mridula G, Manohar P, Chanchalani L, Subhash A. On generalized composite fractional derivative. *Walailak Journal of Science and Technology (WJST)*. 2014; 11(12): 1069-1076.
- [17] Alha S, Mallah IA, Manohar P. On reaction-diffusion equation with generalized composite fractional derivative. *European Journal of Mathematics and Statistics*. 2022; 3(6): 20-26.
- [18] Anumaka MC. Analysis and applications of Laplace/Fourier transformations in electric circuit. *International Journal of Research and Reviews in Applied Sciences (IJRRAS)*. 2012; 12(2): 333-339.
- [19] Rodrigo MR, Mamon RS. An application of Mellin transform techniques to a Black-Scholes equation problem. *Analysis and Applications*. 2007; 5(1): 51-66.
- [20] Mullineux N, Reed JR. On hankel and mellin transforms. *International Journal of Mathematical Education in Science and Technology*. 1978; 9(2): 139-145.
- [21] Elzaki TM. The new integral transform Elzaki transform. *Global Journal of Pure and Applied Mathematics*. 2011; 7(1): 57-64.
- [22] Khalid M, Sultana M, Zaidi F, Arshad U. Application of Elzaki transform method on some fractional differential equations. *Mathematical Theory and Modeling*. 2015; 5(1): 89-96.
- [23] Rashid S, Hammouch Z, Aydi H, Ahmad AG, Alsharif AM. Novel computations of the time-fractional Fisher's model via generalized fractional integral operators by means of the Elzaki transform. *Fractal and Fractional*. 2021; 5(3): 94.
- [24] Sedeeg AKH. A coupling Elzaki transform and homotopy perturbation method for solving nonlinear fractional heat-like equations. *American Journal of Mathematical and Computer Modelling*. 2016; 1(1): 15-20.
- [25] Metzler R, Klafter J. The random walk's guide to anomalous diffusion: a fractional dynamics approach. *Physics Reports*. 2000; 339(1): 1-77.
- [26] Prabhakar TR. A singular integral equation with a generalized Mittag Leffler function in the kernel. *Yokohama Mathematical Journal*. 1971; 19(1): 7-15.
- [27] Kim H. The time shifting theorem and the convolution for Elzaki transform. *International Journal of Pure and Applied Mathematics*. 2013; 87(2): 261-271.
- [28] Mohamed EA. Elzaki transformation for linear fractional differential equations. *Journal of Computational and Theoretical Nanoscience*. 2015; 12(9): 2303-2305.
- [29] Singh Y, Gill V, Kundu S, Kumar D. On the Elzaki transform and its applications in fractional free electron laser equation. *Acta Universitatis Sapientiae, Mathematica*. 2019; 11(2): 419-429.
- [30] Gupta VG, Sharma B. Application of Sumudu transform in reaction-diffusion systems and nonlinear waves. *Applied Mathematical Sciences*. 2010; 4(9-12): 435-446.