

Research Article

New Parametric Polynomials of U -Charlier-Poisson Type: Properties and Szász-Type Operators Including These Polynomials

Cesarano Clemente^{1*}, Alejandro Urieles², Javier Villa², María José Ortega³

¹Section of Mathematics, UniNettuno University, Corso Vittorio Emanuele II, 39, 00186 Rome, Italy

²Mathematics Program, University of the Atlantic, Km 7 Via Pto. Colombia, Barranquilla, Colombia

³Department of Natural and Exact Sciences, University of the Coast, Barranquilla, Colombia

E-mail: c.cesarano@uninettunouniversity.net

Received: 6 June 2025; **Revised:** 16 July 2025; **Accepted:** 24 July 2025

Abstract: In this article, we introduce a new family of parametric U -Charlier-Poisson type polynomials, denoted by $G_n^{[2+J]}(x; \alpha, \beta, \lambda)$. Then, some properties are studied, such as its explicit representation, the orthogonality relationship, and its connection with the derivative of the harmonic function. Subsequently, Szász-type operators are applied to the new family of polynomials to study convergence properties using Korovkin's theorem.

Keywords: Charlier polynomials, Korovkin theorem, Brenke type operators

MSC: 11B68, 11B83, 11B39, 05A19

1. Introduction

Throughout this article, \mathbb{N} will mean the set of natural numbers; $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$, likewise \mathbb{R} , \mathbb{R}^+ , and \mathbb{C} will denote the set of real numbers, positive real numbers, and the set of complex numbers. As usual, will denote by $C[0, \infty)$ the set of all functions f continuous in the interval $[0, \infty)$. The notation $UC[0, \infty)$ will denote the space of functions uniformly continuous on $[0, \infty)$. The space of all polynomials in one variable with real coefficients is denoted by \mathbb{P} , and $\log(z)$ denotes the principal value of the multi-valued logarithm function. In [1], a famous theorem about linear operators is published, known as the Korovkin theorem, which states that a sequence of linear operators under certain conditions converges uniformly in each subset of the locally compact domain. Korovkin theorem, in its many applications, was also used to demonstrate the convergence of Szász operators, which are defined by (see [2, p.239, Eq. (2)]):

$$S_n(f; x) := e^{-nx} \sum_{k=0}^{\infty} \frac{(nx)^k}{k!} f\left(\frac{k}{n}\right), \quad (1)$$

where $f \in C[0, \infty)$, $n \in \mathbb{N}$, and $x \geq 0$. The generalizations of Szász operators by using polynomials, especially defined via generating functions, have been frequently studied lately. These kinds of generalizations provide a range of new sequences

of operators to approximation theory highly advantageous when interpolating positive continuous functions [2]. A known generalization of (1) can be obtained using the Appell polynomials given by (cf. [3]):

$$P_n(f; x) := \frac{e^{-nx}}{g(1)} \sum_{k=0}^{\infty} p_k(nx) f\left(\frac{k}{n}\right), \quad (2)$$

considering that $p_k(x) \geq 0$, for $x \in [0, \infty)$ and $g(1) \neq 0$.

Some time later, Varma et al. ([3, p.122 Eq. (1.7)]), generalized (1) in the following way: first, they use the Brenke-type polynomials, which are defined by the following generating function:

$$\zeta(z)\xi(xz) = \sum_{k=0}^{\infty} p_k(x) \frac{z^k}{k!}, \quad (3)$$

where ζ and ξ are analytical functions. Second, they introduce the positive linear operators including the Brenke-type, polynomials which are given by [3, p.121, Eq. (1.12)]:

$$L_n(f; x) := \frac{1}{\zeta(1)\xi(nx)} \sum_{k=0}^{\infty} p_k(nx) f\left(\frac{k}{n}\right), \quad (4)$$

where $x \geq 0$ and $n \in \mathbb{N}$. It is observed that if $\xi(z) = e^z$ in (3), then the operators (4) concerning (3) lead to (2) with respect to the Appell polynomials, and if $\xi(z) = e^z$ and $\zeta(z) = 1$ in (4), we have (1).

On the other hand, when using the Brenke-type polynomials given in (3), with $\xi(z) = e^z$ and $\zeta(t) = \left(1 - \frac{z}{a}\right)^u$, we have the classic Charlier-Poisson polynomials, which are defined by [4, p.458, Eq. (1.2)]:

$$e^z \left(1 - \frac{z}{a}\right)^u = \sum_{k=0}^{\infty} C_k(a, u) \frac{z^k}{k!}, \quad a \neq 0. \quad (5)$$

Then, Varma et al. introduce the positive linear operators involving Charlier-Poisson polynomials (see [5, p.119, Eq. (1.6)]) by replacing $\xi(z) = e^z$ and $\zeta(z) = \left(1 - \frac{z}{a}\right)^u$ in (4), as follows:

$$L_n(f; x, a) = e^{-1} \left(1 - \frac{1}{a}\right)^{(a-1)nx} \sum_{k=0}^{\infty} \frac{C_k(a, -(a-1)nx)}{k!} f\left(\frac{k}{n}\right), \quad (6)$$

where $a > 1$ and $x \geq 0$. We see that if in (6) we take on both sides $a \rightarrow \infty$ and $x \rightarrow x - \frac{1}{n}$, then we get the Szász operators given in (1). The convergence and bounding properties of these operators were also investigated [5]. Furthermore, in [6], a study of Charlier-Poisson polynomials is presented, in particular, their explicit representation given by

$$C_n(x, \alpha) = \sum_{k=0}^n \binom{n}{k} \binom{x}{k} k! (-\alpha)^{n-k}.$$

The Charlier-Poisson polynomials $C_n(x, \alpha)$, $x \in \mathbb{N}_0$, and, $\alpha > 0$, are orthogonal with respect to the Poisson distribution with mean α , that is,

$$\sum_{x=0}^{\infty} C_m(x, \alpha) C_n(x, \alpha) \frac{e^{-\alpha} \alpha^x}{x!} = \alpha^{-n} n! \delta_{m,n}, \quad m, n \in \mathbb{N}_0,$$

where δ_{mn} is the Kronecker delta. On the topic of polynomial families and their various extensions, a remarkably large amount of research has appeared in the literature (see, for example, [7–13]).

Our contribution aims to introduce a new family of discrete polynomials, called new parametric U -Charlier-Poisson type polynomials, and we investigate some of their properties. Thus, the operators obtained from Brenke-type polynomials are applied to the said polynomials. The outline of this work is as follows: In Section 2, we study some basic results of operators obtained from Brenke-type polynomials applied to Charlier-Poisson polynomials and other results necessary for developing this work. In Section 3, we introduce the new parametric U -Charlier-Poisson type polynomials and explore some of their properties. In Section 4, we investigate the orthogonality relation. Finally, in Section 5, we apply the Szász-type operators (4), obtained from Brenke-type polynomials to the new family of polynomials to study the convergence and bounding properties.

2. Background and previous results

Let f be some function of a real variable x . The backward and forward difference operators Δ and ∇ respectively, are defined as (see [14, p.19-20]):

$$\begin{aligned} \nabla f(x) &:= f(x) - f(x-1), \\ \Delta f(x) &:= f(x+1) - f(x). \end{aligned} \tag{7}$$

Given two real-valued sequence functions $\{a_k(x)\}$ and $\{b_k(x)\}$, if $b_{-1} = 0$, then (see [14, p.20])

$$\sum_{k=0}^{\infty} (\Delta a_k(x)) b_k(x) = - \sum_{k=0}^{\infty} a_k(x) \nabla b_k(x). \tag{8}$$

Furthermore, for $f_1(x)$ and $f_2(x)$ with real values, the following property is satisfied (cf. [15]):

$$\nabla(f_1(x)f_2(x)) = f_1(x)\nabla f_2(x) + f_2(x-1)\nabla f_1(x). \tag{9}$$

The falling factorial x of order n is (see [16])

$$\langle x \rangle_n := x(x-1) \cdots (x-n+1), \quad \text{with } \langle x \rangle_0 = 1, \tag{10}$$

and the rising factorial x of order n is (see [16])

$$(x)_n := x(x+1) \cdots (x+n-1), \quad (x)_0 = 1. \quad (11)$$

The rising factorial and the falling factorial fulfill the following relationship (see [15]):

$$(x)_n = \frac{\Gamma(n+x)}{\Gamma(x)}, \quad (12)$$

$$\langle x \rangle_n = \frac{x!}{(x-n)!}, \quad (13)$$

where $\Gamma(x)$ is the Gamma function.

On the other hand, the digamma function $\psi(x)$ is defined as the logarithmic derivative of the gamma function $\Gamma(x)$ (see [17])

$$\psi(x) = \frac{d}{dx} \log \Gamma(x) = \frac{\Gamma'(x)}{\Gamma(x)}. \quad (14)$$

The generalized harmonic number function is given by (see [17])

$$H_n^m(x) = \sum_{k=0}^{n-1} \frac{1}{(k+x)^m}, \quad n, m \in \mathbb{N}. \quad (15)$$

If $m = 1$ in (15), then

$$H_n^1(x) = \sum_{k=0}^{n-1} \frac{1}{k+x}. \quad (16)$$

If $x = 0$ in (15), we have

$$H_n^m(0) = H_n^m = \sum_{k=1}^n \frac{1}{k^m},$$

where H_n^m denotes the so-called n -th harmonic numbers of order m .

Notice that from (12) and (14) follows

$$\begin{aligned} \frac{d}{dx}(x)_n &= \frac{\Gamma(n+x)}{\Gamma(x)} \left(\frac{d}{dx} \ln(\Gamma(n+x)) - \frac{d}{dx} \ln(\Gamma(x)) \right) \\ &= \frac{\Gamma(n+x)}{\Gamma(x)} (\psi(n+x) - \psi(x)). \end{aligned} \quad (17)$$

By (12), (16), and (17), we obtain

$$(x)_n = \frac{1}{H_n^1(x)} \frac{d}{dx} (x)_n. \quad (18)$$

The Stirling numbers of the first kind $s(n, k)$, appear as the coefficients in the following generating function (see [18]):

$$\frac{(\log(1+z))^k}{k!} = \sum_{n=k}^{\infty} s(n, k) \frac{z^n}{n!}.$$

In addition, they also satisfy

$$\langle x \rangle_n = \sum_{k=0}^n s(n, k) x^k. \quad (19)$$

Note that from (19), we can write

$$(1+z)^x = \sum_{n=0}^{\infty} \binom{x}{n} z^n = \sum_{n=0}^{\infty} \langle x \rangle_n \frac{z^n}{n!} = \sum_{n=0}^{\infty} \left(\sum_{k=0}^n s(n, k) \frac{z^n}{n!} \right) x^k. \quad (20)$$

Now, (cf. [6, p.170 Eq. (1.1)]) it is possible to represent the Charlier-Poisson polynomials given in (5) as follows:

$$e^{-\alpha w} (1+w)^x = \sum_{n=0}^{\infty} (-\alpha)^n C_n(x, \alpha) \frac{w^n}{n!}, \quad (21)$$

with $\alpha \neq 0$. Note that by taking $\alpha = a$, $w = -\frac{z}{a}$, and $x = u$ in (21) we have (5).

It is worth noting that the classical orthogonal polynomials possess a weight function that conforms to the Pearson equation of the form

$$\nabla [\sigma(x) \omega(x)] = \tau(x) \omega(x), \quad (22)$$

with σ a polynomial of degree ≤ 2 and τ a polynomial of degree ≤ 1 . We note that in (22) the backward difference operator ∇ , given in (7), is used for orthogonal polynomials on the lattice and it is replaced by differentiation in the case of orthogonal polynomials on an interval of the real line. The Pearson equation is an important part of the theory of classical orthogonal polynomials because it enables us to derive many useful properties of these polynomials.

Let $f \in UC[0, \infty)$, If $\delta > 0$, the modulus of continuity of the function f , denoted by $\omega(f; \delta)$ is defined by (cf. [5])

$$\omega(f; \delta) := \sup_{x, y \in [0, \infty)} |f(x) - f(y)|, \text{ where } |x - y| < \delta. \quad (23)$$

Additionally, it is well known that,

$$|f(x) - f(y)| \leq \omega(f, \delta) \left(\frac{|x-y|}{\delta} + 1 \right). \quad (24)$$

Also, we have if f is uniformly continuous, then

$$|f(x) - f(y)| \leq \omega(f, \delta). \quad (25)$$

The following Proposition summarizes some properties of the operators defined in (6).

Proposition 1 For $n \in \mathbb{N}$, let $L_n(f; x, a)$ be the positive linear operators involving Charlier-Poisson polynomials in the variable $x \geq 0$. Then, the following statements hold.

1. [5, Lemma 1] The operators defined in (6) satisfy the following identities:

$$(i) L_n(1; x, a) = 1.$$

$$(ii) L_n(s; x, a) = x + \frac{1}{n}.$$

$$(iii) L_n(s^2; x, a) = x^2 + \frac{x}{n} \left(3 + \frac{1}{a-1} \right) + \frac{2}{n^2}.$$

2. [5, Theorem 1] Let E be the set given by

$$S = \{f: [0, \infty) \rightarrow \mathbb{R}: |f(x)| \leq M e^{Ax}, A \in \mathbb{R} \text{ and } M \in \mathbb{R}^+\}.$$

If $f \in C[0, \infty) \cap S$, then

$$\lim_{n \rightarrow \infty} L_n(f; x, a) = f(x).$$

That is, the operators defined in (6) converge uniformly on every compact subset of $[0, \infty)$.

3. [5, Theorem 2] Let $f \in UC[0, \infty) \cap S$. Then the operators L_n given in (6) satisfy

$$|L_n(f; x, a) - f(x)| \leq \left\{ 1 + \sqrt{x \left(1 + \frac{1}{a-1} \right) + \frac{2}{n}} \right\} \omega \left(f; \frac{1}{\sqrt{n}} \right),$$

with ω given by (23).

3. New parametric U -Charlier-Poisson type polynomials and some of their properties

In this section, we shall introduce a new class of discrete polynomials, which we denote by $G_n^{[2+J]}(x; \alpha, \beta, \lambda)$ and will we call new parametric U -Charlier-Poisson type polynomials. Furthermore, we obtain some of their properties.

Definition 1 For a fixed $J \in \mathbb{N}$, $\beta, \lambda \in \mathbb{R}$ and $\alpha \neq 0$, the new family of parametric U -Charlier-Poisson type polynomials $G_n^{[2+J]}(x; \alpha, \beta, \lambda)$ in the variable $x \in \mathbb{R}$ are defined by the means of the power series expansion at 0 of the following generating function:

$$u(x; z; \alpha, \beta, \lambda) = \left[\beta e^{-\alpha z} + \lambda^n (-\alpha)^{-J} \prod_{m=2}^{2+J} \frac{2^m - 1}{2^m - 2} \right] (1+z)^x = \sum_{n=0}^{\infty} G_n^{[2+J]}(x; \alpha, \beta, \lambda) \frac{z^n}{n!}. \quad (26)$$

From (26) and taking $A_j(\lambda, \alpha) = \lambda^n (-\alpha)^{-J} \prod_{m=2}^{2+J} \frac{2^m - 1}{2^m - 2}$, the first parametric U -Charlier-Poisson type polynomials are obtained, which are:

$$G_0^{[2+J]}(x; \alpha, \beta, \lambda) = \beta + A_j(\lambda, \alpha),$$

$$G_1^{[2+J]}(x; \alpha, \beta, \lambda) = -\alpha\beta + A_j(\lambda, \alpha) + x(\beta + A_j(\lambda, \alpha)),$$

$$G_2^{[2+J]}(x; \alpha, \beta, \lambda) = \alpha^2\beta - 2\alpha\beta x + x(x-1)(\beta + A_j(\lambda, \alpha)),$$

$$G_3^{[2+J]}(x; \alpha, \beta, \lambda) = -\alpha^3\beta + \alpha^2\beta x - 2\alpha\beta x(x-1) + x(x-1)(x-2)(-\alpha\beta + \beta + A_j(\lambda, \alpha)).$$

Note that if $\beta = 1$ and $\lambda = 0$, $z = w$ in (26), we have the classic Charlier-Poisson polynomials given in (21). Therefore, the generating function of $G_n^{[2+J]}(x; \alpha, \beta, \lambda)$ in (26) includes, as its special cases, the generating function of the Charlier-Poisson polynomials, i.e., $C_n(x, \alpha) = G_n^{[2+J]}(x; \alpha, 1, 0)$.

Substituting $x = 0$ in (26), we have

$$u(0; z; \alpha, \beta, \lambda) = \left[\beta e^{-\alpha z} + \lambda^n (-\alpha)^{-J} \prod_{m=2}^{2+J} \frac{2^m - 1}{2^m - 2} \right] = \sum_{n=0}^{\infty} G_n^{[2+J]}(0; \alpha, \beta, \lambda) \frac{z^n}{n!}, \quad (27)$$

which leads us to the following definition.

Definition 2 (Parametric U -Charlier-Poisson numbers) The associated sequence of parametric U -Charlier-Poisson numbers is given by evaluating the polynomial at $x = 0$:

$$G_n^{[2+J]}(\alpha, \beta, \lambda) := G_n^{[2+J]}(0; \alpha, \beta, \lambda), \quad n \in \mathbb{N}_0.$$

These numbers arise as coefficients in expansions where the variable x is fixed, and they are central in the structural and algebraic properties studied later.

We can compute a few values of the numbers $G_n^{[2+J]}(\alpha, \beta, \lambda)$ as follows:

$$G_0^{[2+J]}(\alpha, \beta, \lambda) = \beta + A_j(\lambda, \alpha),$$

$$G_3^{[2+J]}(\alpha, \beta, \lambda) = -\beta\alpha^3,$$

$$G_1^{[2+J]}(\alpha, \beta, \lambda) = -\beta\alpha,$$

$$G_4^{[2+J]}(\alpha, \beta, \lambda) = \beta\alpha^4,$$

$$G_2^{[2+J]}(\alpha, \beta, \lambda) = \alpha^2\beta,$$

$$G_5^{[2+J]}(\alpha, \beta, \lambda) = -\beta\alpha^5.$$

We now isolate a sequence of coefficients arising from the parametric structure of the generating function. In particular, the contribution corresponding to the second summand in the generating function

$$\lambda^n (-\alpha)^{-J} \prod_{m=2}^{2+J} \frac{2^m - 1}{2^m - 2} (1+z)^x$$

yields a sequence of numbers $U_n^{[2+J]}(\alpha)$, which we define via the expansion:

$$\lambda^n (-\alpha)^{-J} \prod_{m=2}^{2+J} \frac{2^m - 1}{2^m - 2} (1+z)^x = \sum_{n=0}^{\infty} U_n^{[2+J]}(\alpha) \frac{z^n}{n!}.$$

These coefficients encapsulate the purely parametric contribution independent of β , and their structure is useful in analyzing limiting cases and algebraic identities. Although a closed-form expression is not available, they may be seen as structurally analogous to convolution-type sequences arising in exponential generating functions.

One can use $A_j(\lambda, \alpha)$ in the following manner:

$$A_j(\lambda, \alpha) = \lambda^n (-\alpha)^{-J} \prod_{m=2}^{2+J} \frac{2^m - 1}{2^m - 2} = \sum_{n=0}^{\infty} U_n^{[2+J]}(\alpha) \frac{\lambda^n}{n!}. \quad (28)$$

Whereby some $U_n^{[2+J]}(\alpha)$ are

$$U_0^{[2+J]}(\alpha) = (-\alpha)^{-J} \prod_{m=2}^{2+J} \frac{2^m - 1}{2^m - 2},$$

$$U_2^{[2+J]}(\alpha) = 2(-\alpha)^{-J} \prod_{m=2}^{2+J} \frac{2^m - 1}{2^m - 2},$$

$$U_1^{[2+J]}(\alpha) = (-\alpha)^{-J} \prod_{m=2}^{2+J} \frac{2^m - 1}{2^m - 2},$$

$$U_3^{[2+J]}(\alpha) = 6(-\alpha)^{-J} \prod_{m=2}^{2+J} \frac{2^m - 1}{2^m - 2}.$$

Proposition 2 Let $\beta \in \mathbb{R} - \{0\}$, $J \in \mathbb{N}$ fixed, and $\left\{ G_k^{[2+J]}(\alpha, \beta, \lambda) \right\}_{k=0}^{\infty}$ be a parametric U -Charlier-Poisson type sequence of numbers defined as in (27). Then, the following relationship is fulfilled:

$$G_n^{[2+J]}(\alpha, \beta, \lambda) = (-1)^n \beta \alpha^n, \quad (29)$$

with

$$G_0^{[2+J]}(\alpha, \beta, \lambda) = \beta + A_j(\lambda, \alpha). \quad (30)$$

Proof. By using (27) follows

$$\begin{aligned} \sum_{n=0}^{\infty} G_n^{[2+J]}(\alpha, \beta, \lambda) \frac{z^n}{n!} &= A_j(\lambda, \alpha) + \beta \sum_{n=0}^{\infty} (-1)^n \alpha^n \frac{z^n}{n!} \\ \Leftrightarrow G_0^{[2+J]}(\alpha, \beta, \lambda) + \sum_{n=1}^{\infty} G_n^{[2+J]}(\alpha, \beta, \lambda) \frac{z^n}{n!} &= \beta \sum_{n=0}^{\infty} (-1)^n \alpha^n \frac{z^n}{n!} + A_j(\lambda, \alpha). \end{aligned}$$

With what we have,

$$G_0^{[2+J]}(\alpha, \beta, \lambda) = \sum_{n=1}^{\infty} \left[(-1)^n \alpha^n \beta - G_n^{[2+J]}(\alpha, \beta, \lambda) \right] \frac{z^n}{n!} + (\beta + A_j(\lambda, \alpha)). \quad (31)$$

From (31) follows (29) and (30). Proposition 2 is proved. \square

With its proof, the following proposition provides a concise formula for the parametric U -Charlier-Poisson type polynomials $G_n^{[2+J]}(x; \alpha, \beta, \lambda)$.

Proposition 3 Given a fixed $J \in \mathbb{N}$, let $\left\{ G_n^{[2+J]}(x; \alpha, \beta, \lambda) \right\}_{n=0}^{\infty}$ be a parametric U -Charlier-Poisson type sequence of polynomials, defined as in (26). Then, the following explicit representation hold:

$$G_n^{[2+J]}(x; \alpha, \beta, \lambda) = \beta (-\alpha)^n C_n(x, \alpha) + \lambda^n (-\alpha)^{-J} \left[\prod_{m=2}^{2+J} \frac{2^m - 1}{2^m - 2} \right] \langle x \rangle_n, \quad (32)$$

where $\langle x \rangle_n$, is the falling factorial defined in (10).

Proof. Using the generating function of the parametric U -Charlier-Poisson type polynomials given in (26), we have

$$\begin{aligned} \sum_{n=0}^{\infty} G_n^{[2+J]}(x; \alpha, \beta, \lambda) \frac{z^n}{n!} &= \left[\beta e^{-\alpha z} + \lambda^n (-\alpha)^{-J} \prod_{m=2}^{2+J} \frac{2^m - 1}{2^m - 2} \right] (1+z)^x \\ &= \beta \sum_{n=0}^{\infty} (-\alpha)^n C_n(x, \alpha) \frac{z^n}{n!} + \lambda^n (-\alpha)^{-J} \left[\prod_{m=2}^{2+J} \frac{2^m - 1}{2^m - 2} \right] \sum_{n=0}^{\infty} \binom{x}{n} z^n \\ &= \beta \sum_{n=0}^{\infty} (-\alpha)^n C_n(x, \alpha) \frac{z^n}{n!} + \lambda^n (-\alpha)^{-J} \left[\prod_{m=2}^{2+J} \frac{2^m - 1}{2^m - 2} \right] \sum_{n=0}^{\infty} \langle x \rangle_n \frac{z^n}{n!} \\ &= \sum_{n=0}^{\infty} \left[\beta (-\alpha)^n C_n(x, \alpha) + \lambda^n (-\alpha)^{-J} \prod_{m=2}^{2+J} \frac{2^m - 1}{2^m - 2} \langle x \rangle_n \right] \frac{z^n}{n!}. \end{aligned}$$

Considering the above expression, we thus have (32). Proposition 3 is demonstrated. \square

Proposition 4 For a fixed $J \in \mathbb{N}$, let $\left\{G_k^{[2+J]}(x; \alpha, \beta, \lambda)\right\}_{k=0}^{\infty}$ be a parametric U -Charlier-Poisson type sequence of polynomials defined by (26). If $\beta \rightarrow 0$ and $\lambda \rightarrow 1$, then the following identity holds:

$$\sum_{k=0}^n \binom{n}{k} G_k^{[2+J]}(x; \alpha, 0, 1) \alpha^{n-k} C_{n-k}(-\alpha, -x) = \alpha^n \sum_{n=0}^{\infty} n! \prod_{m=2}^{2+J} \frac{2^m - 1}{2^m - 2}. \quad (33)$$

Proof. Let us write (26) as

$$\begin{aligned} [\beta e^{-\alpha z}(1+z)^x + A_j(\lambda, \alpha)(1+z)^x] e^{\alpha z}(1+z)^{-x} &= \left(\sum_{n=0}^{\infty} G_n^{[2+J]}(x; \beta, \alpha, \lambda) \frac{z^n}{n!} \right) e^{\alpha z}(1+z)^{-x} \\ &= \left(\sum_{n=0}^{\infty} G_n^{[2+J]}(x; \beta, \alpha, \lambda) \frac{z^n}{n!} \right) \left(\sum_{n=0}^{\infty} \alpha^n C_n(-\alpha, -x) \frac{z^n}{n!} \right). \end{aligned}$$

From the above expression and (28), we have

$$\begin{aligned} \beta + e^{\alpha z} A_j(\lambda, \alpha) &= \sum_{n=0}^{\infty} \sum_{k=0}^n \binom{n}{k} G_k^{[2+J]}(x; \beta, \alpha, \lambda) \alpha^{n-k} C_{n-k}(-\alpha, -x) \frac{z^n}{n!}. \\ \Leftrightarrow \beta + \sum_{n=0}^{\infty} \alpha^n \left(\sum_{n=0}^{\infty} U_n^{[2+J]}(\lambda; \alpha) \frac{\lambda^n}{n!} \right) \frac{z^n}{n!} &= \sum_{n=0}^{\infty} \sum_{k=0}^n \binom{n}{k} G_k^{[2+J]}(x; \beta, \alpha, \lambda) \alpha^{n-k} C_{n-k}(-\alpha, -x) \frac{z^n}{n!}. \end{aligned}$$

Then, taking $\beta \rightarrow 0$ and $\lambda \rightarrow 1$, follows

$$\sum_{n=0}^{\infty} \alpha^n \left(\sum_{n=0}^{\infty} n! \prod_{m=2}^{2+J} \frac{2^m - 1}{2^m - 2} \right) \frac{z^n}{n!} = \sum_{n=0}^{\infty} \sum_{k=0}^n \binom{n}{k} G_k^{[2+J]}(x; \beta, \alpha, \lambda) \alpha^{n-k} C_{n-k}(-\alpha, -x) \frac{z^n}{n!},$$

from which (33) follows. Proposition 4 is demonstrated. \square

Proposition 5 For a fixed $J \in \mathbb{N}$, $\beta \in \mathbb{R} - \{0\}$, let $\left\{G_k^{[2+J]}(x; \alpha, \beta, \lambda)\right\}_{k=0}^{\infty}$ be a parametric U -Charlier-Poisson type sequence of polynomials defined by (26). Then, we have the following relationship:

$$\alpha^n A_j(\lambda, \alpha) C_n(x, -\alpha) + \sum_{k=0}^n \beta s(n, k) x^k = \sum_{l=0}^n \binom{n}{l} G_l^{[2+J]}(x; \alpha, \beta, \lambda) \alpha^{n-l}, \quad (34)$$

where $s(n, k)$ is defined by (19).

Proof. From (26), implies that

$$\begin{aligned}\beta(1+z)^x &= \left[\sum_{n=0}^{\infty} G_n^{[2+J]}(x; \alpha, \beta, \lambda) \frac{z^n}{n!} - A_j(\lambda, \alpha)(1+z)^x \right] \frac{1}{e^{-\alpha z}} \\ &= \left(\sum_{n=0}^{\infty} G_n^{[2+J]}(x; \alpha, \beta, \lambda) \frac{z^n}{n!} \right) \left(\sum_{n=0}^{\infty} \alpha^n \frac{z^n}{n!} \right) - A_j(\lambda, \alpha) \sum_{n=0}^{\infty} \alpha^n C_n(x, -\alpha) \frac{z^n}{n!}.\end{aligned}$$

Now, using (20) follows:

$$\begin{aligned}\beta \sum_{n=0}^{\infty} \left(\sum_{k=0}^n s(n, k) x^k \right) \frac{z^n}{n!} &= \sum_{n=0}^{\infty} \sum_{l=0}^n \binom{n}{l} G_l^{[2+J]}(x; \alpha, \beta, \lambda) \alpha^{n-l} \frac{z^n}{n!} - A_j(\lambda, \alpha) \sum_{n=0}^{\infty} \alpha^n C_n(x, -\alpha) \frac{z^n}{n!} \\ &= \sum_{n=0}^{\infty} \left(\sum_{l=0}^n \binom{n}{l} G_l^{[2+J]}(x; \alpha, \beta, \lambda) \alpha^{n-l} - A_j(\lambda, \alpha) \alpha^n C_n(x, -\alpha) \right) \frac{z^n}{n!},\end{aligned}$$

which yields (34). Our Proposition 5 is proven. \square

Proposition 6 For a fixed $J \in \mathbb{N}$, $\beta \in \mathbb{R} - \{0\}$ the following relations hold for the parametric U -Charlier-Poisson type polynomials defined by (26):

$$n \frac{\partial}{\partial x} G_{n-1}^{[2+J]}(x; \alpha, \beta, \lambda) = \sum_{k=1}^n (-1)^{k-1} (n-k) \langle n \rangle_k G_{n-k-1}^{[2+J]}(x; \alpha, \beta, \lambda), \quad (G_{-n}^{[2+J]} \equiv 0), \quad (35)$$

$$\frac{1}{\alpha} G_{n+1}^{[2+J]}(x; \alpha, \beta, \lambda) - \mathfrak{K}(x; z; \alpha) \frac{\partial}{\partial x} G_n^{[2+J]}(x; \alpha, \beta, \lambda) + G_n^{[2+J]}(x; \alpha, \beta, \lambda) - A_j(\lambda, \alpha) \langle x \rangle_n = 0, \quad (36)$$

where $\alpha \in \mathbb{R} - \{0\}$, $z \in \mathbb{C} - \{0, -1\}$, $n \in \mathbb{N}$ with

$$\mathfrak{K}(x; z; \alpha) = \frac{x}{\alpha} \left[\frac{1}{(1+z) \log(1+z)} \right], \quad (37)$$

and $A_j(\lambda, \alpha)$ given in (28).

Proof. To prove (35), we note that by differentiating (26) with respect to x , we can write

$$\begin{aligned}\sum_{n=0}^{\infty} \frac{\partial}{\partial x} G_n^{[2+J]}(x; \alpha, \beta, \lambda) \frac{z^n}{n!} &= \left(\sum_{n=1}^{\infty} (-1)^n \frac{z^n}{n} \right) \left(\sum_{n=0}^{\infty} G_n^{[2+J]}(x; \alpha, \beta, \lambda) \frac{z^n}{n!} \right) \\ \Leftrightarrow \sum_{n=1}^{\infty} \frac{\partial}{\partial x} G_{n-1}^{[2+J]}(x; \alpha, \beta, \lambda) \frac{z^{n-1}}{(n-1)!} &= \left(\sum_{n=1}^{\infty} (-1)^n (n-1)! \frac{z^n}{n} \right) \left(\sum_{n=0}^{\infty} G_{n-1}^{[2+J]}(x; \alpha, \beta, \lambda) \frac{z^{n-1}}{(n-1)!} \right)\end{aligned}$$

$$\Leftrightarrow \sum_{n=1}^{\infty} n \frac{\partial}{\partial x} G_{n-1}^{[2+J]}(x; \alpha, \beta, \lambda) \frac{z^n}{n!} = \sum_{n=1}^{\infty} \sum_{k=1}^n (-1)^{k-1} (k-1)! \binom{n}{k} (n-k) G_{n-1-k}^{[2+J]}(x; \alpha, \beta, \lambda) \frac{z^n}{n!}$$

$$\Leftrightarrow n \frac{\partial}{\partial x} G_{n-1}^{[2+J]}(x; \alpha, \beta, \lambda) = \sum_{k=1}^n (-1)^{k-1} (k-1)! \binom{n}{k} (n-k) G_{n-1-k}^{[2+J]}(x; \alpha, \beta, \lambda).$$

Of the above expression and applying (13) follows (35).

Now to prove (36), we derive (26) with respect to z as follows:

$$\frac{\partial}{\partial z} u(x; z; \alpha, \beta, \lambda) = \sum_{n=0}^{\infty} G_{n+1}^{[2+J]}(x; \alpha, \beta, \lambda) \frac{z^n}{n!}, \quad (38)$$

and

$$\frac{\partial}{\partial z} u(x; z; \alpha, \beta, \lambda) = \frac{x}{(1+z)} [(1+z)^x (\beta e^{-\alpha z} + A_j(\lambda, \alpha))] - \alpha \beta e^{-\alpha z} (1+z)^x. \quad (39)$$

Likewise, if we derive (26) with respect to x , we have the following:

$$\frac{\partial}{\partial x} u(x; z; \alpha, \beta, \lambda) = \sum_{n=0}^{\infty} \frac{\partial}{\partial x} G_n^{[2+J]}(x; \alpha, \beta, \lambda) \frac{z^n}{n!}, \quad (40)$$

$$\frac{\partial}{\partial x} u(x; z; \alpha, \beta, \lambda) = (1+z)^x \log(1+z) (\beta e^{-\alpha z} + A_j(\lambda, \alpha)). \quad (41)$$

By using (38), (39), (40), and (41), we obtain

$$\begin{aligned} & \frac{1}{\alpha} \frac{\partial}{\partial z} u(x; z; \alpha, \beta, \lambda) - \frac{1}{\alpha} \left[\frac{x}{(1+z) \log(1+z)} \right] \frac{\partial}{\partial x} u(x; z; \alpha, \beta, \lambda) \\ & + u(x; z; \alpha, \beta, \lambda) - (1+z)^x A_j(\lambda, \alpha) = 0 \\ \Leftrightarrow & \frac{1}{\alpha} \sum_{n=0}^{\infty} G_{n+1}^{[2+J]}(x; \alpha, \beta, \lambda) \frac{z^n}{n!} - \frac{1}{\alpha} \sum_{n=0}^{\infty} \left[\frac{x}{(1+z) \log(1+z)} \right] \frac{\partial}{\partial x} G_n^{[2+J]}(x; \alpha, \beta, \lambda) \frac{z^n}{n!} \\ & + \sum_{n=0}^{\infty} G_n^{[2+J]}(x; \alpha, \beta, \lambda) \frac{z^n}{n!} - A_j(\lambda, \alpha) \sum_{n=0}^{\infty} \binom{x}{n} z^n = 0 \end{aligned}$$

$$\begin{aligned} &\Leftrightarrow \sum_{n=0}^{\infty} \frac{1}{\alpha} G_{n+1}^{[2+J]}(x; \alpha, \beta, \lambda) \frac{z^n}{n!} - \sum_{n=0}^{\infty} \frac{1}{\alpha} \left[\frac{x}{(1+z) \log(1+z)} \right] \frac{\partial}{\partial x} G_n^{[2+J]}(x; \alpha, \beta, \lambda) \frac{z^n}{n!} \\ &+ \sum_{n=0}^{\infty} G_n^{[2+J]}(x; \alpha, \beta, \lambda) \frac{z^n}{n!} - \sum_{n=0}^{\infty} A_j(\lambda, \alpha) \langle x \rangle_n \frac{z^n}{n!} = 0. \end{aligned}$$

Of the previous equation taking $\mathfrak{K}(x; z; \alpha)$ as in (37), (36) follows. Proposition 6 is proved. \square

Proposition 7 For a fixed $J \in \mathbb{N}$, $\beta \in \mathbb{R} - \{0\}$, let $G_n^{[2+J]}(x; \alpha, \beta, \lambda)$ be the parametric U -Charlier-Poisson type polynomials. Then the following statement holds:

$$\frac{d}{dx}(x)_n = \frac{H_n(x)}{\beta} \left[-\lambda^n (-\alpha)^{-J} \prod_{m=2}^{2+J} \frac{2^m - 1}{2^m - 2} + \sum_{k=0}^n (-1)^n \alpha^k G_{n-k}^{[2+J]}(-x; \alpha, \beta, \lambda) \right], \quad (42)$$

using $(x)_n$ given by (11) and $H_n(x) = H_n^1(x)$ given in (16).

Proof. Taking $z \rightarrow -z$ and $x \rightarrow -x$ in (26), we have

$$\begin{aligned} \beta(1-z)^{-x} &= e^{-\alpha z} \sum_{n=0}^{\infty} (-1)^n G_n^{[2+J]}(-x; \alpha, \beta, \lambda) \frac{z^n}{n!} - A_j(\lambda, \alpha) e^{-\alpha z} (1-z)^{-x} \\ &= \left(\sum_{n=0}^{\infty} (-1)^n \alpha^n \frac{z^n}{n!} \right) \left(\sum_{n=0}^{\infty} (-1)^n G_n^{[2+J]}(-x; \alpha, \beta, \lambda) \frac{z^n}{n!} \right) - A_j(\lambda, \alpha) \sum_{n=0}^{\infty} (-1)^n (-\alpha)^n C_n(-x, \alpha) \frac{z^n}{n!} \\ &= \sum_{n=0}^{\infty} \sum_{k=0}^n \binom{n}{k} (-1)^n \alpha^k G_{n-k}^{[2+J]}(-x; \alpha, \beta, \lambda) \frac{z^n}{n!} - A_j(\lambda, \alpha) \sum_{n=0}^{\infty} (-1)^n \alpha^n C_n(-x, \alpha) \frac{z^n}{n!}. \end{aligned}$$

Then, for $|z| < 1$, using the Binomial Theorem, we have

$$\begin{aligned} \frac{1}{(1-z)^x} &= \sum_{n=0}^{\infty} \beta^{-1} \left[\sum_{k=0}^n \binom{n}{k} (-1)^n \alpha^k G_{n-k}^{[2+J]}(-x; \alpha, \beta, \lambda) - A_j(\lambda, \alpha) (-1)^n \alpha^n C_n(-x, \alpha) \right] \frac{z^n}{n!} \\ \sum_{n=0}^{\infty} (x)_n \frac{z^n}{n!} &= \sum_{n=0}^{\infty} \beta^{-1} \left[\sum_{k=0}^n \binom{n}{k} (-1)^n \alpha^k G_{n-k}^{[2+J]}(-x; \alpha, \beta, \lambda) - A_j(\lambda, \alpha) (-1)^n \alpha^n C_n(-x, \alpha) \right] \frac{z^n}{n!} \\ (x)_n &= \sum_{k=0}^n (-1)^n \beta^{-1} \alpha^k G_{n-k}^{[2+J]}(-x; \alpha, \beta, \lambda) - \beta^{-1} A_j(\lambda, \alpha) (-1)^n \alpha^n C_n(-x, \alpha). \end{aligned}$$

This way, using (18) follows (42). This completes the demonstration of Proposition 7. \square

4. Orthogonality relationship of the polynomials $G_n^{[2+J]}(x; \alpha, \beta, \lambda)$

The main aim of this section is to obtain the relation of orthogonality satisfied by the polynomials $G_n^{[2+J]}(x; \alpha, \beta, \lambda)$, and apply it to study a relationship between these polynomials and the operator ∇ given in (7).

For a fixed $J \in \mathbb{N}$, we define the parametric U -Charlier-Poisson discrete weight function ω^α as

$$\omega^\alpha(x; \beta, \lambda) = \frac{e^{-\alpha} \alpha^x}{x!} |M(\beta, \lambda, \alpha) + i\Theta(\beta, \lambda, \alpha)|^{-2}, \quad (43)$$

where $\alpha \in \mathbb{R}$, $\alpha < 0$; $\beta, \lambda \in \mathbb{R} - \{0\}$, on the lattice \mathbb{N} ; $z, w \in \mathbb{C}$; $z = a_1 + ia_2$, $w = c_1 + ic_2$ in the circle $C(0, |\eta|)$ and $|\eta| = \min\{|z|, |w|\}$. While $M(\beta, \lambda, \alpha)$ and $\Theta(\beta, \lambda, \alpha)$ are given by

$$\begin{aligned} M(\beta, \lambda, \alpha) &= \beta(\beta + A_j(\lambda, \alpha)(\varepsilon_2 \cos(c_2 \alpha) + \varepsilon_1 \cos(a_2 \alpha))) \\ &\quad + [A_j(\lambda, \alpha)]^2 \varepsilon_1 \varepsilon_2 \cos(\alpha(a_2 + c_2)), \end{aligned} \quad (44)$$

$$\begin{aligned} \Theta(\beta, \lambda, \alpha) &= \beta A_j(\lambda, \alpha)(\varepsilon_2 \sin(c_2 \alpha) + \varepsilon_1 \sin(a_2 \alpha)) \\ &\quad + [A_j(\lambda, \alpha)]^2 \varepsilon_1 \varepsilon_2 \sin(\alpha(a_2 + c_2)), \end{aligned} \quad (45)$$

where $A_j(\lambda, \alpha)$ given in (28), $\varepsilon_1 = e^{a_1 \alpha}$, and $\varepsilon_2 = e^{c_1 \alpha}$.

With the weight $\omega^\alpha(x; \beta, \lambda)$ given in (43), we can introduce on \mathbb{P} the inner product as follows:

$$\langle f, g \rangle_{\omega^\alpha} = \sum_{x=0}^{\infty} f(x)g(x)\omega^\alpha(x; \beta, \lambda),$$

where $f, g \in \mathbb{P}$. Which has positive weights for every $\alpha < 0$.

The Pearson equation concerning (22) for weight (43) is now of the form

$$\nabla \omega^\beta(x; \alpha, \beta, \lambda) = \left(\frac{\alpha - x}{\alpha} \right) \omega^\beta(x; \alpha, \beta, \lambda). \quad (46)$$

Theorem 1 For a fixed $J \in \mathbb{N}$, if $\alpha \in \mathbb{R}$, $\alpha < 0$, $\beta, \lambda \in \mathbb{R} - \{0\}$, and $m, n \in \mathbb{N}_\mu$. Then, the parametric U -Charlier-Poisson type polynomials for the weight (43) satisfy the following orthogonality relation:

$$\sum_{x=0}^{\infty} G_m^{[2+J]}(x; \alpha, \beta, \lambda) G_n^{[2+J]}(x; \alpha, \beta, \lambda) \frac{e^{-\alpha} \alpha^x}{x!} |\Omega(\beta, \lambda, \alpha)|^{-2} = \frac{\Gamma(n+1) \alpha^n}{\Omega(\beta, \lambda, \alpha)} \delta_{m,n}. \quad (47)$$

Where $\Omega(\beta, \lambda, \alpha) = M(\beta, \lambda, \alpha) + i\Theta(\beta, \lambda, \alpha)$.

Proof. One can see that from (26) follows:

$$\begin{aligned}
U(x; z, \alpha, \beta, \lambda) &= \beta \left(\sum_{n=0}^{\infty} (-\alpha)^n \frac{z^n}{n!} \right) \sum_{n=0}^{\infty} \binom{x}{n} z^n + A_j(\lambda, \alpha) \sum_{n=0}^{\infty} \binom{x}{n} z^n \\
&= \beta \sum_{n=0}^{\infty} \sum_{k=0}^n \binom{n}{k} (-\alpha)^{n-k} \langle x \rangle_k \frac{z^n}{n!} + A_j(\lambda, \alpha) \sum_{n=0}^{\infty} \langle x \rangle_n \frac{z^n}{n!} \\
&= \sum_{n=0}^{\infty} \left[\beta \sum_{k=0}^n \binom{n}{k} (-\alpha)^{n-k} \langle x \rangle_k + A_j(\lambda, \alpha) \langle x \rangle_n \right] \frac{z^n}{n!} \\
&= \sum_{n=0}^{\infty} \left[\sum_{k=0}^n \beta \binom{n}{k} (-\alpha)^{n-k} \frac{\langle x \rangle_k}{n!} + A_j(\lambda, \alpha) \frac{\langle x \rangle_n}{n!} \right] z^n.
\end{aligned}$$

This implies that

$$U(x; z, \alpha, \beta, \lambda) = \sum_{n=0}^{\infty} D_n^{[2+J]}(x; \alpha, \beta, \lambda) z^n, \quad (48)$$

where

$$D_n^{[2+J]}(x; \alpha, \beta, \lambda) = \sum_{k=0}^n \beta \binom{n}{k} (-\alpha)^{n-k} \frac{\langle x \rangle_k}{n!} + A_j(\lambda, \alpha) \frac{\langle x \rangle_n}{n!}.$$

Similarly, we obtain

$$D_m^{[2+J]}(x; \alpha, \beta, \lambda) = \sum_{k=0}^m \beta \binom{m}{k} (-\alpha)^{m-k} \frac{\langle x \rangle_k}{m!} + A_j(\lambda, \alpha) \frac{\langle x \rangle_m}{m!}.$$

On the other hand, we have

$$\begin{aligned}
U(x; z, \alpha, \beta, \lambda) U(x; w, \alpha, \beta, \lambda) &= [\beta e^{-\alpha z} + A_j(\lambda, \alpha)] [\beta e^{-\alpha w} + A_j(\lambda, \alpha)] (1+z)^x (1+w)^x \\
&= e^{-\alpha z} e^{-\alpha w} (\beta + A_j(\lambda, \alpha) e^{\alpha z}) (\beta + A_j(\lambda, \alpha) e^{\alpha w}) (1+z)^x (1+w)^x,
\end{aligned}$$

and so, we have that

$$\begin{aligned}
\sum_{k=0}^{\infty} \frac{\alpha^k U(x; z, \alpha, \beta, \lambda) U(x; w, \alpha, \beta, \lambda)}{k!} &= (\beta + A_j(\lambda, \alpha) e^{\alpha z})(\beta + A_j(\lambda, \alpha) e^{\alpha w}) e^{-\alpha z} e^{-\alpha w} e^{\alpha(1+z)(1+w)} \\
&= (\beta + A_j(\lambda, \alpha) e^{\alpha z})(\beta + A_j(\lambda, \alpha) e^{\alpha w}) e^{\alpha} e^{\alpha z w}.
\end{aligned}$$

So,

$$\sum_{k=0}^{\infty} U(x; z, \alpha, \beta, \lambda) U(x; w, \alpha, \beta, \lambda) \frac{e^{-\alpha} \alpha^k}{k!} = (\beta + A_j(\lambda, \alpha) e^{\alpha z})(\beta + A_j(\lambda, \alpha) e^{\alpha w}) \sum_{n=0}^{\infty} \alpha^n \frac{z^n w^n}{n!}. \quad (49)$$

Applying equation (48) to the left-hand side of (49) yields

$$\begin{aligned}
\sum_{k=0}^{\infty} U(x; z, \alpha, \beta, \lambda) U(x; w, \alpha, \beta, \lambda) \frac{e^{-\alpha} \alpha^k}{k!} &= \sum_{k=0}^{\infty} \frac{e^{-\alpha} \alpha^k}{k!} \sum_{n=0}^{\infty} D_n^{[2+J]}(x; \alpha, \beta, \lambda) z^n \sum_{m=0}^{\infty} D_m^{[2+J]}(x; \alpha, \beta, \lambda) w^m \\
&= \sum_{m,n=0}^{\infty} \sum_{k=0}^{\infty} D_m^{[2+J]}(x; \alpha, \beta, \lambda) D_n^{[2+J]}(x; \alpha, \beta, \lambda) \frac{e^{-\alpha} \alpha^k}{k!} z^n w^m. \quad (50)
\end{aligned}$$

By combining Equation (49) with Equation (50), we have that

$$\sum_{n=0}^{\infty} (\beta + e^{\alpha z} b)(\beta + e^{\alpha w} b) \frac{\alpha^n z^n w^n}{n!} = \sum_{m,n=0}^{\infty} \sum_{k=0}^{\infty} D_m^{[2+J]}(x; \alpha, \beta, \lambda) D_n^{[2+J]}(x; \alpha, \beta, \lambda) \frac{e^{-\alpha} \alpha^k}{k!} z^n w^m.$$

Which results in

$$\begin{aligned}
\sum_{k=0}^{\infty} D_m^{[2+J]}(k; \alpha, \beta, \lambda) D_n^{[2+J]}(k; \alpha, \beta, \lambda) \frac{e^{-\alpha} \alpha^k}{k!} &= \begin{cases} \left[\frac{\alpha^n (\beta + A_j(\lambda, \alpha) e^{\alpha z})(\beta + A_j(\lambda, \alpha) e^{\alpha w})}{n!} \right], & \text{if } m = n, \\ 0, & \text{if } m \neq n. \end{cases} \\
\Leftrightarrow \sum_{k=0}^{\infty} D_m^{[2+J]}(k; \alpha, \beta, \lambda) D_n^{[2+J]}(k; \alpha, \beta, \lambda) \frac{e^{-\alpha} \alpha^k}{k!} &= \left[\frac{\alpha^n (\beta + A_j(\lambda, \alpha) e^{\alpha z})(\beta + A_j(\lambda, \alpha) e^{\alpha w})}{n!} \right] \delta_{m,n}.
\end{aligned}$$

And so we can see that

$$\sum_{k=0}^{\infty} G_m^{[2+J]}(x, \alpha, \beta, \lambda) G_n^{[2+J]}(x, \alpha, \beta, \lambda) \frac{e^{-\alpha} \alpha^k}{k!} = n! \alpha^n (\beta + A_j(\lambda, \alpha) e^{\alpha z})(\beta + A_j(\lambda, \alpha) e^{\alpha w}) \delta_{m,n}. \quad (51)$$

Now, from Equation (51), we consider the following product:

$$\begin{aligned}
(\beta + A_j(\lambda, \alpha)e^{\alpha z})(\beta + A_j(\lambda, \alpha)e^{\alpha w}) &= \beta^2 + \beta \varepsilon_2 A_j(\lambda, \alpha)e^{ic_2 \alpha} + \beta \varepsilon_1 A_j(\lambda, \alpha)e^{ia_2 \alpha} \\
&+ [A_j(\lambda, \alpha)]^2 \varepsilon_1 \varepsilon_2 e^{ia_2 \alpha} e^{ic_2 \alpha}.
\end{aligned} \tag{52}$$

Finally, we take into consideration the following: we develop the calculations in (52) and substitute Equations (44) and (45) into the result, then organizing (51) with these calculations we get (47), which completes the proof of Theorem 1. \square

Through renewed invocation of (52) with well-defined parameters, we establish the following result.

Corollary 1 For a fixed $J \in \mathbb{N}$, if $\alpha \in \mathbb{R}$, $\alpha < 0$, $\beta, \lambda \in \mathbb{R} - \{0\}$, and $m, n \in \mathbb{N}_\neq$. Assume that $z_1 = a_1 + ia_2$, $z_2 = c_1 + ic_2$, with $a_1, c_1 \rightarrow 0$ and $a_2 \rightarrow c_1 = \zeta$ in the circle $C(0, |\eta|)$. Then, the parametric U -Charlier-Poisson type polynomials satisfy the following orthogonality relation

$$\sum_{x=0}^{\infty} G_m^{[2+J]}(x, \alpha, \beta, \lambda) G_n^{[2+J]}(x, \alpha, \beta, \lambda) \frac{e^{-\alpha} \alpha^x}{x!} |\Omega_1(\beta, \lambda, \alpha)|^{-2} = \frac{\Gamma(n+1) \alpha^n}{\Omega_1(\beta, \lambda, \alpha)} \delta_{m,n}.$$

With $\Omega_1(\beta, \lambda, \alpha) = M_1(\beta, \lambda, \alpha) + i\Theta_1(\beta, \lambda, \alpha)$. Also $M_1(\beta, \lambda, \alpha)$ and $\Theta_1(\beta, \lambda, \alpha)$ are given by

$$M_1(\beta, \lambda, \alpha) = \beta \left(\beta + 2A_j(\lambda, \alpha) \cos(\zeta \alpha) + [A_j(\lambda, \alpha)]^2 \cos(2\alpha \zeta) \right),$$

$$\Theta_1(\beta, \lambda, \alpha) = 2\beta A_j(\lambda, \alpha) \sin(\zeta \alpha) + [A_j(\lambda, \alpha)]^2 \sin(2\zeta \alpha).$$

Using the orthogonality property of the polynomials $G_n^{[2+J]}(x, \alpha, \beta, \lambda)$, the summation by parts given in (8), and the Pearson equation given in (46), we can see the following structure relation:

Proposition 8 The parametric U -Charlier-Poisson type polynomials given in (26), satisfy

$$\Delta G_n^{[2+J]}(x; \alpha, \beta, \lambda) = a_{n-1,n}^\alpha G_{n-1}^{[2+J]}(x; \alpha, \beta, \lambda), \tag{53}$$

where $a_{n-1,n}^\alpha$ are the Fourier coefficients.

Proof. Writing the polynomials $\Delta G_n^{[2+J]}(x; \alpha, \beta, \lambda) = G_n^{[2+J]}(x+1; \alpha, \beta, \lambda) - G_n^{[2+J]}(x; \alpha, \beta, \lambda)$ in terms of $\{G_n^{[2+J]}(x, \alpha, \beta, \lambda)\}_{n \geq 0}$, we have

$$G_n^{[2+J]}(x+1; \alpha, \beta, \lambda) - G_n^{[2+J]}(x; \alpha, \beta, \lambda) = \sum_{k=0}^{n-1} a_{k,n}^\alpha G_k^{[2+J]}(x; \alpha, \beta, \lambda),$$

where

$$a_{k,n}^\alpha = \frac{\left\langle \Delta G_n^{[2+J]}(x; \alpha, \beta, \lambda), G_k^{[2+J]}(x; \alpha, \beta, \lambda) \right\rangle_{\omega^\alpha}}{\left\langle G_k^{[2+J]}(x; \alpha, \beta, \lambda), G_k^{[2+J]}(x; \alpha, \beta, \lambda) \right\rangle_{\omega^\alpha}}, \quad k = 0, 1, \dots, n-1.$$

This way, applying (8) and (9) follows

$$\begin{aligned} & \left\langle G_k^{[2+J]}, G_k^{[2+J]} \right\rangle_{\omega^\alpha} a_{k,n}^\alpha \\ &= \sum_{L=0}^{\infty} \left(\Delta G_n^{[2+J]}(L; \alpha, \beta, \lambda) G_k^{[2+J]}(L; \alpha, \beta, \lambda) \right) \omega^\alpha(L, \beta, \lambda) \\ &= - \sum_{L=0}^{\infty} G_n^{[2+J]}(L; \alpha, \beta, \lambda) \nabla \left(\omega^\alpha(L, \beta, \lambda) G_k^{[2+J]}(L; \alpha, \beta, \lambda) \right) \\ &= - \sum_{L=0}^{\infty} G_n^{[2+J]}(L; \alpha, \beta, \lambda) \left[\omega^\alpha(L, \beta, \lambda) \nabla G_k^{[2+J]}(L; \alpha, \beta, \lambda) + G_k^{[2+J]}(L-1; \alpha, \beta, \lambda) \nabla \omega^\alpha(L) \right] \\ &= - \sum_{L=0}^{\infty} G_n^{[2+J]}(L; \alpha, \beta, \lambda) \omega^\alpha(L, \beta, \lambda) \nabla G_k^{[2+J]}(L; \alpha, \beta, \lambda) \\ &\quad - \sum_{L=0}^{\infty} G_n^{[2+J]}(L-1; \alpha, \beta, \lambda) G_k^{[2+J]}(L-1; \alpha, \beta, \lambda) \nabla \omega^\alpha(L; \beta, \lambda). \end{aligned}$$

Now, due to orthogonality of $G_n^{[2+J]}(L-1; \alpha, \beta, \lambda)$, and since ∇G_k^α has degree $k-1$, we have that the first sum is zero. For the second sum, substituting (46) yields

$$\begin{aligned} \left\langle G_k^{[2+J]}, G_k^{[2+J]} \right\rangle_{\omega^\alpha} a_{k,n}^\alpha &= - \sum_{L=0}^{\infty} G_n^{[2+J]}(L; \alpha, \beta, \lambda) G_k^{[2+J]}(L-1; \alpha, \beta, \lambda) \omega^\alpha(L, \beta, \lambda) \frac{\alpha-L}{\alpha} \\ &= - \frac{1}{\alpha} \sum_{L=0}^{\infty} G_n^{[2+J]}(L; \alpha, \beta, \lambda) G_k^{[2+J]}(L-1; \alpha, \beta, \lambda) \omega^\alpha(L, \beta, \lambda). \end{aligned} \quad (54)$$

This sum is zero for $k+1 < n$, so only $a_{n-1,n}^\alpha$ can be non-zero. Therefore, from (54) follows (53). The proposition 8 is proved. \square

5. Szász-type operators including the parametric U -Charlier-Poisson type polynomials

In this section, we present a positive linear Szász-type operator given by (4) involving the U -Charlier-Poisson type polynomials. With the help of the Korovkin theorem, we study the convergence and some properties.

Definition 3 We define the Szász-type operators, including the generating function of the parametric U -Charlier-Poisson type polynomials given in (26), with $\alpha = a$, $z = -\frac{1}{a}$, and $x = -(a-1)nx$ as follows:

$$J_n(f, x) = (\beta e + A_j(\lambda, \alpha))^{-1} \left(1 - \frac{1}{a}\right)^{(a-1)nx} \sum_{k=0}^{\infty} \frac{G_k^{[2+J]}(-(a-1)nx, a, \beta, \lambda)}{k!} f\left(\frac{k}{n}\right), \quad (55)$$

where $f \in C[0, \infty)$, $n \in \mathbb{N}$, $\beta e \neq A_j(\lambda, \alpha)$, and $x \geq 0$.

Lemma 1 For $n \in \mathbb{N}$ and $x \geq 0$, the operators J_n defined by (55) satisfy the following identities:

1. $J_n(1, x) = 1$,
2. $J_n(s, x) = x + \frac{\beta e}{n(\beta e + A_j(\lambda, \alpha))}$,
3. $J_n(s^2, x) = x^2 + x \left(\frac{1}{n(a-1) + \frac{2\beta e}{n(\beta e + A_j(\lambda, \alpha))} + \frac{1}{n}} \right) + \frac{2\beta e}{n^2(\beta e + A_j(\lambda, \alpha))}$, with $\beta e \neq A_j(\lambda, \alpha)$.

Proof. Using the generating function of the parametric U -Charlier-Poisson type polynomials, given by (26), we can see that

$$\sum_{k=0}^{\infty} \frac{G_k^{[2+J]}(-(a-1)nx, a, \beta, \lambda)}{k!} = \left(1 - \frac{1}{a}\right)^{-(a-1)nx} (\beta e + A_j(\lambda, \alpha)), \quad (56)$$

$$\sum_{k=0}^{\infty} \frac{k G_k^{[2+J]}(-(a-1)nx, a, \beta, \lambda)}{k!} = \left(1 - \frac{1}{a}\right)^{-(a-1)nx} [\beta e + nx(\beta e + A_j(\lambda, \alpha))], \quad (57)$$

$$\begin{aligned} \sum_{k=0}^{\infty} \frac{k^2 G_k^{[2+J]}(-(a-1)nx, a, \beta, \lambda)}{k!} &= \left(1 - \frac{1}{a}\right)^{-(a-1)nx} \\ &\times [n^2 x^2 (\beta e + A_j(\lambda, \alpha)) + n^2 x (\beta e + A_j(\lambda, \alpha)) \Phi + 2\beta e], \end{aligned} \quad (58)$$

where

$$\Phi = \left(\frac{1}{n(a-1) + \frac{2\beta e}{n(\beta e + A_j(\lambda, \alpha))} + \frac{1}{n}} \right).$$

Then, multiplying each equation in (56)-(58) by their respective right multiplicative inverses and applying the operator definition (55) yields the lemma's assertions. \square

Theorem 2 Let $S = \{f: [0, \infty) \rightarrow \mathbb{R}: |f(x)| \leq Me^{Ax}\}$, where $A \in \mathbb{R}$. If $f \in C[0, \infty) \cap S$, then

$$\lim_{n \rightarrow \infty} J_n(f, x) = f.$$

That is, the operators defined in (55) converge uniformly on every compact subset of $[0, \infty)$.

Proof. By using Lemma 1, we have

$$\lim_{n \rightarrow \infty} J_n(s^i; x, a) = x^i, \quad i = 0, 1, 2.$$

In this way, using Korovkin's Theorem [19], convergence is guaranteed in each compact subset of $[0, \infty)$. □

The next result gives the rate of convergence of the sequence J_n to f by means of the modulus of continuity.

Theorem 3 Let $f \in UC[0, \infty) \cap S$. Then the operators J_n satisfy the inequality that follows:

$$|J_n(f, x) - f(x)| \leq \left\{1 + \sqrt{\Upsilon_n(x; \beta, \lambda)}\right\} \omega\left(f; \frac{1}{\sqrt{n}}\right),$$

with

$$\Upsilon_n(x; \beta, \lambda) = \left[\frac{x}{n} \left[\frac{n^2(\beta e + H)^2 - 2\beta e}{n^3(a-1)(\beta e + H)^2 + 3\beta e + H} \right] + \frac{2\beta e}{n^2(\beta e + H)} \right], \quad a \neq 1$$

where $H = \left(\beta e + \lambda(-\alpha)^{-J} \prod_{m=2}^{2+J} \frac{2^m - 1}{2^m - 2} \right)^{-1}$.

Proof. By using (23), (25), the Definition of the new operators given in (55), and identity 1 of Lemma 1, we can write

$$|J_n(f, x) - f(x)| = \left| H \left(1 - \frac{1}{a}\right)^{(a-1)nx} \sum_{k=0}^{\infty} \frac{G_k^{[2+J]}(-(a-1)nx, a, \beta, \lambda)}{k!} f\left(\frac{k}{n}\right) - 1 \cdot f(x) \right|.$$

Thereupon

$$\begin{aligned} |J_n(f, x) - f(x)| &= \left| H \left(1 - \frac{1}{a}\right)^{(a-1)nx} \sum_{k=0}^{\infty} \frac{G_k^{[2+J]}(-(a-1)nx, a, \beta, \lambda)}{k!} \left(f\left(\frac{k}{n}\right) - f(x)\right) \right| \\ &\leq H \left(1 - \frac{1}{a}\right)^{(a-1)nx} \sum_{k=0}^{\infty} \frac{G_k^{[2+J]}(-(a-1)nx, a, \beta, \lambda)}{k!} \left|f\left(\frac{k}{n}\right) - f(x)\right|. \end{aligned}$$

This way of (24) follows:

$$\begin{aligned}
|J_n(f, x) - f(x)| &\leq \left\{ H \left(1 - \frac{1}{a} \right)^{(a-1)nx} \sum_{k=0}^{\infty} \frac{G_k^{[2+J]}(-(a-1)nx, a, \beta, \lambda)}{k!} \left(\frac{1}{\delta} \left| \frac{k}{n} - x \right| + 1 \right) \omega(f, \delta) \right\} \\
&\leq \left\{ 1 + \frac{1}{\delta} H \left(1 - \frac{1}{a} \right)^{(a-1)nx} \sum_{k=0}^{\infty} \frac{G_k^{[2+J]}(-(a-1)nx, a, \beta, \lambda)}{k!} \left| \frac{k}{n} - x \right| \right\} \omega(f, \delta) \quad (59)
\end{aligned}$$

On the other hand, it holds by Cauchy-Schwarz inequality for series, and Lemma 1 the following:

$$\begin{aligned}
\sum_{k=0}^{\infty} \frac{G_k^{[2+J]}(-(a-1)nx, a, \beta, \lambda)}{k!} \left| \frac{k}{n} - x \right| &\leq \left(H^{-1} \left(1 - \frac{1}{a} \right)^{-(a-1)nx} \right)^{1/2} \\
&\quad \times \left(\sum_{k=0}^{\infty} \frac{G_k^{[2+J]}(-(a-1)nx, a, \beta, \lambda)}{k!} \left(\frac{k}{n} - x \right)^2 \right)^{1/2}.
\end{aligned}$$

Then, taking

$$\phi = \left(H^{-1} \left(1 - \frac{1}{a} \right)^{-(a-1)nx} \right)^{1/2} \left(\sum_{k=0}^{\infty} \frac{G_k^{[2+J]}(-(a-1)nx, a, \beta, \lambda)}{k!} \left(\frac{k}{n} - x \right)^2 \right)^{1/2},$$

is fulfilled

$$\begin{aligned}
\phi &= \left(H^{-1} \left(1 - \frac{1}{a} \right)^{-(a-1)nx} \right)^{1/2} \\
&\quad \times \left(H^{-1} \left(1 - \frac{1}{a} \right)^{-(a-1)nx} H \left(1 - \frac{1}{a} \right)^{(a-1)nx} \sum_{k=0}^{\infty} \frac{G_k^{[2+J]}(-(a-1)nx, a, \beta, \lambda)}{k!} \left(\frac{k}{n} - x \right)^2 \right)^{1/2} \\
&= H^{-1} \left(1 - \frac{1}{a} \right)^{-(a-1)nx} \left[H \left(1 - \frac{1}{a} \right)^{(a-1)nx} \sum_{k=0}^{\infty} \frac{G_k^{[2+J]}(-(a-1)nx, a, \beta, \lambda)}{k!} \left(\frac{k^2}{n^2} - 2\frac{kx}{n} + x^2 \right) \right]^{1/2} \\
&= H^{-1} \left(1 - \frac{1}{a} \right)^{-(a-1)nx} \left[J_n(s^2, x) - 2xJ_n(s, x) + x^2J_n(1, x) \right]^{1/2}.
\end{aligned}$$

So, of (59) and the above expression, we get

$$|J_n(f, x) - f(x)| \leq \left\{ 1 + \frac{1}{\delta} \sqrt{\Upsilon_n(x; \beta, \lambda)} \right\} \omega(f, \delta).$$

□

By choosing $\delta := \delta_n = \frac{1}{\sqrt{n}}$, we arrive at the desired result. Theorem 3 is proved.

Lemma 2 For $x \in [0, \infty)$, the sequence of operators J_n given in (55) satisfy the following property

$$J_n((s-x)^2, x) = x \left(\frac{1}{n(a-1) + \frac{2\beta e}{n(\beta e + A_j(\lambda, \alpha))} + \frac{1}{n}} - \frac{2\beta e}{n(\beta e + A_j(\lambda, \alpha))} \right) + \frac{2\beta e}{n^2(\beta e + A_j(\lambda, \alpha))},$$

with $\beta e \neq A_j(\lambda, \alpha)$.

Proof. By taking advantage of the linearity property of J_n operators, we have

$$J_n((s-x)^2, x) = J_n(s^2, x) - 2xJ_n(s, x) - x^2J_n(1, x).$$

Next, we apply Lemma 1, we obtain the desired outcome.

□

Theorem 4 Let $f \in C[0, \infty) \cap \mathcal{S}$ and $x \in [0, \infty)$. The operators J_n satisfy the inequality that follows:

$$|J_n(f, x) - f(x)| \leq 2\omega(f; \sqrt{\tau_n}), \quad (60)$$

where

$$\tau_n = x \left(\frac{1}{n(a-1) + \frac{2\beta e}{n(\beta e + A_j(\lambda, \alpha))} + \frac{1}{n}} - \frac{2\beta e}{n(\beta e + A_j(\lambda, \alpha))} \right) + \frac{2\beta e}{n^2(\beta e + A_j(\lambda, \alpha))}.$$

Proof. By the identity 1 of Lemma 1, and using the modulus of continuity property, it is fulfilled

$$\begin{aligned} |J_n(f, x) - f(x)| &\leq H \left(1 - \frac{1}{a} \right)^{(a-1)nx} \sum_{k=0}^{\infty} \frac{G_k^{[2+J]}(-(a-1)nx, a, \beta, \lambda)}{k!} \left| f\left(\frac{k}{n}\right) - f(x) \right| \\ &\leq \left\{ 1 + H \left(1 - \frac{1}{a} \right)^{(a-1)nx} \frac{1}{\delta} \sum_{k=0}^{\infty} \frac{G_k^{[2+J]}(-(a-1)nx, a, \beta, \lambda)}{k!} \left| \frac{k}{n} - x \right| \right\} \omega(f, \delta). \end{aligned}$$

On the other hand, by Lemma 2 and the Cauchy-Schwarz inequality, it holds

$$\begin{aligned}
\sum_{k=0}^{\infty} \frac{G_k^{[2+J]}(-(a-1)nx, a, \beta, \lambda)}{k!} \left| \frac{k}{n} - x \right| &\leq \sqrt{(H)^{-1} \left(1 - \frac{1}{a}\right)^{(a-1)nx}} \\
&\times \left\{ \sum_{k=0}^{\infty} \frac{G_k^{[2+J]}(-(a-1)nx, a, \beta, \lambda)}{k!} \left| \frac{k}{n} - x \right| \right\}^{1/2} \\
&\leq H \left(1 - \frac{1}{a}\right)^{-(a-1)nx} \{\tau_n\}^{1/2}.
\end{aligned}$$

This way, $|J_n(f, x) - f(x)| \leq \left\{1 + \frac{1}{\delta} \sqrt{\tau_n}\right\}^{1/2}$. Thus, by taking $\delta = \sqrt{\tau_n}$, we have the desired result. \square

Data availability

Data sharing is not applicable to this article as no datasets were generated or analysed during the current study.

Conflict of interest

The authors declare no competing financial interest.

References

- [1] Korovkin P. On convergence of linear positive operators in the space of continuous functions (Russia). *Doklady Academy of Sciences*. 1953; 90: 961-964.
- [2] Szasz O. Generalization of S. Bernstein's polynomials to the infinite interval. *Journal of Research of the National Bureau of Standards*. 1950; 45: 239-245.
- [3] Varma S, Sezgin S, Gurhan I. Generalization of Szasz operators involving Brenke type polynomials. *Computers and Mathematics with Applications*. 2012; 64: 121-127. Available from: <https://doi.org/10.1016/j.camwa.2012.01.025>.
- [4] Ozmen N, Erkus DE. On the Poisson-Charlier polynomials. *Serdica Mathematical Journal*. 2015; 41: 458-470.
- [5] Varma S, Tasdelen F. Szasz type operators involving Charlier polynomials. *Mathematical and Computer Modelling*. 2012; 56: 118-122. Available from: <https://doi.org/10.1016/j.mcm.2011.12.017>.
- [6] Chihara TS. *An Introduction to Orthogonal Polynomials*. New York: Gordon and Breach; 1978.
- [7] Castilla L, Cesarano C, Bedoya D, Ramírez W, Agarwal P, Jain S. A generalization of the Apostol-X type Frobenius-Genocchi polynomials of level ι . In: Agarwal P, Cattani C, Momani S. (eds.) *Fractional Differential Equations: Theoretical Aspects and Applications, Advanced Studies in Complex Systems*. Academic Press; 2024. p.11-26. Available from: <https://doi.org/10.1016/B978-0-44-315423-2.00007-2>.
- [8] Dattoli G, Cesarano C. On a new family of Hermite polynomials associated to parabolic cylinder functions. *Applied Mathematics and Computation*. 2003; 141(1): 143-149. Available from: [https://doi.org/10.1016/S0096-3003\(02\)00328-4](https://doi.org/10.1016/S0096-3003(02)00328-4).
- [9] Dattoli G, Ricci PE, Cesarano C. The Lagrange polynomials, the associated generalizations, and the umbral calculus. *Integral Transforms and Special Functions*. 2003; 14(2): 181-186. Available from: <https://doi.org/10.1080/1065246031000098186>.
- [10] Dattoli G, Lorenzutta S, Cesarano C. Bernstein polynomials and operational methods. *Journal of Computational Analysis and Applications*. 2006; 8(4): 369-377.

- [11] Urieles A, Ramírez W, Ortega MJ, Bedoya D. Fourier expansion and integral representation for generalized Apostol-type Frobenius-Euler polynomials. *Advances in Difference Equations*. 2020; 2020: 534. Available from: <https://doi.org/10.1186/s13662-020-02988-0>.
- [12] Tabinda N, Shahid AW, Parvez A, Ramírez W. Introducing a novel family of Δ_h -Sheffer polynomials and their interconnected hybrid variants. *Journal of Mathematics and Computer Science*. 2024; 36(3): 317-325. Available from: <https://doi.org/10.22436/jmcs.036.03.06>.
- [13] Mohra Z, Shahid AW, Ramírez W, Mohammad A, Fuentes F. Properties and applications of Bell polynomials of two variables. *Journal of Mathematics and Computer Science*. 2024; 35(3): 291-303. Available from: <https://doi.org/10.22436/jmcs.035.03.03>.
- [14] Nikiforov AF, Suslov SK, Uvarov VB. *Classical Orthogonal Polynomials of a Discrete Variable*. Berlin Heidelberg: Springer-Verlag; 1991.
- [15] Gradshteyn IS. *Table of Integrals, Series and Products*. New York: Gordon and Breach; 1978.
- [16] Koekoek R, Lesky PA, Swarttouw RF. *Hypergeometric Orthogonal Polynomials and Their q -Analogues*. Berlin Heidelberg: Springer-Verlag; 2010.
- [17] Olaikhan A. *An Introduction to the Harmonic Series and Logarithmic Integrals*. Ali Shadhar Olaikhan; 2021.
- [18] Srivastava HM, Choi J. *Zeta and q -Zeta Functions and Associated Series and Integrals*. London: Elsevier; 2012. Available from: <https://dx.doi.org/10.1016/C2010-0-67023-4>.
- [19] Altomare F, Campiti M. *Korovkin-Type Approximation Theory and Its Applications, vol. 17*. Berlin, Germany: Walter De Gruyter; 1994.