

Research Article

Novel Results on Applications of Differential Subordination on Analytic Functions Associated with Zeta-Riemann Fractional Differential Operator

Ekram E. Ali^{1*}, Rabha M. El-ashwah², Abeer H. Alblowy¹, Altaf Alshuhail¹, Fozaiyah A. Alhubairah¹

¹Department of Mathematics, College of Science, University of Hail, Hail, 2440, Saudi Arabia

²Department of Mathematics, Faculty of Science, Damietta University, New Damietta, 34517, Egypt
E-mail: e.ahmad@uoh.edu.sa

Received: 30 June 2025; **Revised:** 21 October 2025; **Accepted:** 5 November 2025

Abstract: The findings of this study are connected with geometric function theory and are obtained by using subordination-based techniques in conjunction with Zeta-Riemann fractional differential operator information. We used the Zeta-Riemann fractional differential operator to investigate a certain classes of analytic functions. It is shown that, for particular choice of parameters for the new generalized classes, the classes of starlike, close-to-convex, and α -convex functions emerges. Many classes of univalent functions are studied using convolution and subordination principle. Furthermore, some classes are defined and developed by using the Zeta-Riemann fractional differential operator. The connections between the classes are presented in the definitions and associated remarks, and characterization properties are also proved, including combinations of functions belonging to those classes and inclusion relations.

Keywords: analytic function, differential subordination, fractional operator

MSC: 30C45, 30C80

1. Introduction and preliminaries

Geometric Functions Theory (GFT) is an amalgamation of geometry along with analysis. The theory of analytic univalent and multivalent functions are two of the most important aspects of the GFT. Investigations in these areas constitute an ancient topic in mathematics, especially in Complex analysis, that has drawn quite a few scholars due to the absolute elegance of their geometrical properties as well as numerous study opportunities. Inarguably, the most significant field of complex analysis to feed just one or many variables is the investigation of univalent functions. Univalent functions are covered in detail in the standard works of Duren [1] and Goodman [2]. The idea of analytic function subordination was first introduced by Littlewood [3], and Rogosinski [4] developed the phrase and established the fundamental results utilizing subordination. The concept of subordination was recently employed by Srivastava and Owa [5] to explore a number of fascinating properties of the generalized hypergeometric function. A generalization of differential inequalities, known as differential subordinations, was the subject of an article by Miller and Mocanu [6]. There have been numerous fascinating properties of GFT that were studied and investigated by several authors, see [7–16].

Let $\mathcal{A}$ denote the class of function satisfying $f(0) = f'(0) - 1 = 0$ written as:

$$f(\xi) = \xi + \sum_{\kappa=2}^{\infty} a_{\kappa} \xi^{\kappa}, \quad (1)$$

which are analytic and univalent in the open unit disc $\mathcal{U} = \{\xi : |\xi| < 1\}$. If f and g are analytic in $\mathcal{U}$, f is subordinate to g , denoted $f(\xi) \prec g(\xi)$, if there exists an analytic function ϖ , with $\varpi(0) = 0$ and $|\varpi(\xi)| < 1$ for all $\xi \in \mathcal{U}$, such that $f(\xi) = g(\varpi(\xi))$, $\xi \in \mathcal{U}$. If the function g is univalent in $\mathcal{U}$, $f(\xi) \prec g(\xi)$ is given as (see [17–19]):

$$f(0) = g(0) \text{ and } f(\mathcal{U}) \subset g(\mathcal{U}).$$

For two functions $f_{\iota}(\xi) \in \mathcal{A}$ ($\iota = 1, 2$) are given by

$$f_{\iota}(\xi) = \xi + \sum_{\kappa=2}^{\infty} a_{\kappa, \iota} \xi^{\kappa},$$

we define the convolution of $f_1(\xi)$ and $f_2(\xi)$ as (see, for example [20], p.246)

$$(f_1 * f_2)(\xi) = \xi + \sum_{\kappa=2}^{\infty} a_{\kappa, 1} a_{\kappa, 2} \xi^{\kappa} = (f_2 * f_1)(\xi).$$

The Zeta-Riemann fractional differential operator is an intriguing extension of classical differential operators, derived from the fusion of the ideas embedded in the Riemann zeta function and the principles of fractional calculus. It emerges from the need to analyze non-local phenomena in mathematical physics and applied mathematics through the lens of fractional derivatives. The fractional calculus, as a mathematical field, explores derivatives and integrals of arbitrary orders, allowing for a more nuanced understanding of dynamical systems that exhibit non-integer order behavior. By incorporating the zeta function, which encodes rich information about number theory and complex analysis, the Zeta-Riemann fractional differential operator captures additional complexity and characteristics from both domains, creating a powerful analytical tool.

Tayyab and Atshan [21] investigate the following fractional differential operator

$$\mathfrak{D}_{(s, k, w)}^{\alpha} f(\xi) = \sum_{\kappa=1}^{\infty} \frac{(w+1)^{\frac{s(\alpha-1)+1}{k}} \Gamma\left(\frac{\kappa}{w+1} + 1\right) \Gamma\left(\frac{\delta + \eta + 1}{2}\right)}{k^{\frac{s(\alpha-1)+1}{k}} \Gamma\left(\frac{\kappa}{w+1} - \frac{s(\alpha-1)+1}{k} + 1\right)} a_{\kappa} \xi^{(w+1)(1 - \frac{s(\alpha-1)+1}{k}) + \kappa - 1}, \quad (2)$$

$$\left(s \in \mathbb{N}, k > 0, w \geq 0, 0 \leq \frac{s(\alpha-1)+1}{k} < 1\right).$$

In [22] the Hurwitz-Lerch Zeta function is defined as following

$$\Phi(\xi, s, \alpha) = \sum_{\kappa=0}^{\infty} \frac{\xi^{\kappa}}{(\kappa + \alpha)^s}, \quad (3)$$

$s \in \mathbb{C}, \alpha \in \mathbb{C} \setminus \mathbb{Z}^-, \operatorname{Re}(s) > 1$, when $|\xi| = 1$.

Several interesting properties and characteristics of the above defined Hurwitz-Lerch Zeta function was studied and investigated by several authors see [23–26].

We define the new Hadamard product fractional differential operator.

$$\begin{aligned} \mathfrak{D}_s^{(\alpha, \beta)} f(\xi) &= \left[D_{(s, 1, 0)}^{\alpha} f(\xi) \right] * \left[\xi^{-(s(\alpha-1)+1)} (\Phi(\xi, s-1, \alpha+\beta) - \alpha^{-s}) \right] \\ &= \sum_{\kappa=1}^{\infty} \frac{\kappa!}{\Gamma(\kappa - s(\alpha-1)) (\kappa + \alpha + \beta)^{s-1}} a_{\kappa} \xi^{\kappa - [s(\alpha-1)+1]}. \end{aligned} \quad (4)$$

We note that if $s = 1$, then we have Srivastava fractional differential operator in [27] as:

$$\mathfrak{D}_1^{(\alpha, \beta)} f(\xi) = \sum_{\kappa=1}^{\infty} \frac{\kappa!}{\Gamma(\kappa - \alpha + 1)} a_{\kappa} \xi^{\kappa - \alpha}. \quad (5)$$

Shexo et al. [28] introduced a new operator (Zeta-Riemann Fractional differential operator) $\mathscr{D}_s^{(\alpha, \beta)} : \mathcal{A} \rightarrow \mathcal{A}$ as follows:

$$\begin{aligned} \mathscr{D}_s^{(\alpha, \beta)} f(\xi) &= (1 + \alpha + \beta)^{s-1} \Gamma(1 - s(\alpha-1)) \xi^{[s(\alpha-1)+1]} \mathfrak{D}_s^{(\alpha, \beta)} f(\xi) \\ &= \xi + \sum_{\kappa=2}^{\infty} \frac{\Gamma(1 - s(\alpha-1)) (1 + \alpha + \beta)^{s-1} \kappa!}{\Gamma(\kappa - s(\alpha-1)) (\kappa + \alpha + \beta)^{s-1}} a_{\kappa} \xi^{\kappa}. \end{aligned} \quad (6)$$

The operator $\mathscr{D}_s^{(\alpha, \beta)} f(\xi)$ is linear and bounded in the unit disk $\mathcal{U}$ and satisfying

$$\xi \left(\mathscr{D}_s^{(\alpha, \beta)} f(\xi) \right)' = s(1 - \alpha) \mathscr{D}_s^{(\alpha + \frac{1}{s}, \beta - \frac{1}{s})} f(\xi) - [s(1 - \alpha) - 1] \mathscr{D}_s^{(\alpha, \beta)} f(\xi). \quad (7)$$

The study of the Zeta-Riemann fractional differential operator is ongoing, with researchers continually uncovering its implications in both pure and applied mathematics. By delving into properties such as stability, existence of solutions, and numerical approximations, mathematicians and scientists alike can leverage this operator to develop innovative models that reflect the intricate dynamics of real-world systems. The interplay between fractional calculus and the analytical properties of the Riemann zeta function also paves the way for future investigations into connections with other areas of mathematics, including chaos theory, fractals, and complex networks.

The primary objective of this paper is to study the dependence concepts with an additional subset for univalent functions alongside a different operator which uses subordination-based techniques in conjunction with Zeta-Riemann fractional differential operator information. We will introduce many classes of univalent functions by using the

convolution and subordination principle. Further, we develop some classes by using the Hurwitz-Lerch Zeta operator and we illustrate the connections between the classes in the definition or in associated remarks and characterization properties are also, we prove including combinations of functions belonging to those classes and inclusion relations.

Let Ψ be the class of analytic univalent convex functions in $\mathcal{U}$, with $\mu(0) = 1$ and $\operatorname{Re}\mu(\xi) > 0$.

The next lemmas are used to demonstrate our results:

Lemma 1 [29] Let $\beta, \sigma \in \mathbb{C}$. Also let $\mu \in \mathcal{A}$ be convex univalent in $\mathcal{U}$ with $\operatorname{Re}[\beta\mu(\xi) + \sigma] > 0$ ($\xi \in \mathcal{U}$), $\mu(0) = 1$ and $p(\xi) \in \mathcal{A}$ with $p(\xi) = 1 + p_1\xi + p_2\xi^2 + \dots$ is analytic in $\mathcal{U}$. If

$$p(\xi) + \frac{\xi p'(\xi)}{\beta p(\xi) + \sigma} \prec \mu(\xi) \quad (\xi \in \mathcal{U})$$

then $p(\xi) \prec \mu(\xi)$.

Lemma 2 [30] Let $\beta, \sigma \in \mathbb{C}$. Also suppose $\mu \in \mathcal{A}$ be convex univalent in $\mathcal{U}$ with $\mu(0) = 1$ and $\operatorname{Re}[\beta\mu(\xi) + \sigma] > 0$ ($\xi \in \mathcal{U}$), and $q(\xi) \in \mathcal{A}$ with $q(0) = 1$ and $q(\xi) \prec \mu(\xi)$ ($\xi \in \mathcal{U}$). If $p(\xi) = 1 + p_1\xi + p_2\xi^2 + \dots$ is analytic in $\mathcal{U}$,

$$p(\xi) + \frac{\xi p'(\xi)}{\beta q(\xi) + \sigma} \prec \mu(\xi) \quad (\xi \in \mathcal{U}),$$

then

$$p(\xi) \prec \mu(\xi).$$

Bu using the operator $\mathcal{D}_s^{(\alpha, \beta)} f(\xi)$, we investigate the class $\mathfrak{R}_s^{(\alpha, \beta)}(\vartheta; \mu)$, of analytic functions $f = \{f_1, f_2, \dots, f_\vartheta\}$ on open unit disc $\mathcal{U}$ satisfying $\frac{\xi (\mathcal{D}_s^{(\alpha, \beta)} f_i(\xi))'}{\frac{1}{\vartheta} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} f_j(\xi)} \prec \mu(\xi)$ ($f_i \in \mathcal{A}$, $i = 1, 2, \dots, \vartheta$, $\xi \in \mathcal{U}$), where $\xi^{-1} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} f_j(\xi) \neq 0$

and μ is convex univalent in $\mathcal{U}$ with $\mu(0) = 1$. Also we define $F = \{F_1, F_2, \dots, F_\vartheta\}$ where $F_i(\xi) = \frac{\varsigma + 1}{\xi \varsigma} \int_0^\xi t^{\varsigma-1} f_i(t) dt$ ($\varsigma > 0$; $i = 1, 2, \dots, \vartheta$) and proved that $F \in \mathfrak{R}_s^{(\alpha, \beta)}(\vartheta; \mu)$, whenever $f \in \mathfrak{R}_s^{(\alpha, \beta)}(\vartheta; \mu)$. Additional classes of this type are indicated by $\mathfrak{C}_s^{(\alpha, \beta)}(\vartheta; \mu)$, $\mathfrak{P}_s^{(\alpha, \beta)}(\vartheta; \delta, \mu)$ and $\mathfrak{R}_s^{(\alpha, \beta)}(\vartheta; \delta, \mu)$ are presented and examined here using convolutions with subordination approach.

2. Main results

Throughout this paper, unless otherwise mentioned, we set $s \geq 1$, $\alpha, \beta \in \mathbb{R}^+$ and $\vartheta \in \mathbb{N}$.

2.1 The class $\mathfrak{R}_s^{(\alpha, \beta)}(\vartheta; \mu)$

Definition 1 Let $f = \{f_1, f_2, \dots, f_\vartheta\}$, $f_i \in \mathcal{A}$, $1 \leq i \leq \vartheta$ be such that

$$\frac{\xi (\mathcal{D}_s^{(\alpha, \beta)} f_i(\xi))'}{\frac{1}{\vartheta} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} f_j(\xi)} \prec \mu(\xi) \quad (\xi \in \mathcal{U}; i = 1, 2, \dots, \vartheta),$$

where $\xi^{-1} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} f_j(\xi) \neq 0$ in $\mathcal{U}$, μ is convex univalent in $\mathcal{U}$ with $\mu(0) = 1$. Then we say that $f = \{f_1, f_2, \dots, f_{\vartheta}\}$ belongs to the class $\mathfrak{R}_s^{(\alpha, \beta)}(\vartheta; \mu)$.

Remark 1 (i) $\mathfrak{R}_1^{(0, 0)}\left(1; \frac{1-\xi}{1+\xi}\right) = S^*$, the class of starlike functions introduced by Robertson [31];

(ii) $\mathfrak{R}_1^{(0, 0)}\left(1; \frac{1+A\xi}{1+B\xi}\right) = S^*(A, B)$, $(-1 \leq B < A \leq 1)$, this class investigate by Janowski [32];

(iii) For $\vartheta = 1$ and $\mu(\xi) = \frac{1-\xi}{1+\xi}$, we have

$$f(\xi) = \xi + \frac{\Gamma(\kappa - s(\alpha - 1))(\kappa + \alpha + \beta)^{s-1}}{\kappa\Gamma(1 - s(\alpha - 1))(1 + \alpha + \beta)^{s-1}\kappa!} \xi^{\kappa} \in \mathfrak{R}_s^{(\alpha, \beta)}\left(1; \frac{1-\xi}{1+\xi}\right) (\kappa \geq 2).$$

Theorem 1 Let $f = \{f_1, f_2, \dots, f_{\vartheta}\} \in \mathfrak{R}_s^{(\alpha, \beta)}(\vartheta; \mu)$ and $F(\xi) = \frac{1}{\vartheta} \sum_{i=1}^{\vartheta} f_i(\xi)$. Then $F(\xi)$ satisfies the condition

$$\frac{\xi \left(\mathcal{D}_s^{(\alpha, \beta)} F(\xi) \right)'}{\mathcal{D}_s^{(\alpha, \beta)} F(\xi)} \prec \mu(\xi) (\xi \in \mathcal{U}), \quad (8)$$

where $\xi^{-1} \mathcal{D}_s^{(\alpha, \beta)} F(\xi) \neq 0$.

Proof. Let $f = \{f_1, f_2, \dots, f_{\vartheta}\} \in \mathfrak{R}_s^{(\alpha, \beta)}(\vartheta; \mu)$. Then for any $\xi_0 \in \mathcal{U}$, we have

$$\frac{\xi_0 \left(\mathcal{D}_s^{(\alpha, \beta)} f_i(\xi_0) \right)'}{\frac{1}{\vartheta} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} f_j(\xi_0)} \prec \mu(\mathcal{U})$$

where $\xi_0^{-1} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} f_j(\xi_0) \neq 0$ and hence equals to $\mu(w_i)$ (say) for some $w_i \in \mathcal{U}$, $i = 1, 2, \dots, \vartheta$. So

$$\frac{\sum_{i=1}^{\vartheta} \xi_0 \left(\mathcal{D}_s^{(\alpha, \beta)} f_i(\xi_0) \right)'}{\frac{1}{\vartheta} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} f_j(\xi_0)} = \sum_{i=1}^{\vartheta} \mu(w_i).$$

Let $f(\xi) = \xi + \sum_{k=2}^{\infty} a_k \xi^k$. Then, from (6), we see that

$$\begin{aligned} \mathcal{D}_s^{(\alpha, \beta)} f(\xi) &= f(\xi) * \left\{ \xi + \sum_{\kappa=2}^{\infty} \frac{\Gamma(1 - s(\alpha - 1))(1 + \alpha + \beta)^{s-1}\kappa!}{\Gamma(\kappa - s(\alpha - 1))(\kappa + \alpha + \beta)^{s-1}} \xi^{\kappa} \right\} \\ &= \left(f * \mathfrak{R}_s^{(\alpha, \beta)} \right)(\xi), \end{aligned}$$

where

$$\mathfrak{K}_s^{(\alpha, \beta)} = \xi + \sum_{\kappa=2}^{\infty} \frac{\Gamma(1-s(\alpha-1))(1+\alpha+\beta)^{s-1}\kappa!}{\Gamma(\kappa-s(\alpha-1))(\kappa+\alpha+\beta)^{s-1}} \xi^{\kappa}. \quad (9)$$

Hence

$$\frac{\xi_0 \left(\mathcal{D}_s^{(\alpha, \beta)} F(\xi_0) \right)'}{\mathcal{D}_s^{(\alpha, \beta)} F(\xi_0)} = \frac{\xi_0 \left[\mathfrak{K}_s^{(\alpha, \beta)}(\xi) * \sum_{i=1}^{\vartheta} f_i(\xi_0) \right]'}{\mathfrak{K}_s^{(\alpha, \beta)}(\xi) * \sum_{j=1}^{\vartheta} f_j(\xi_0)}.$$

Since

$$\mathcal{D}_s^{(\alpha, \beta)} \sum_{j=1}^{\vartheta} f_j(\xi) = \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} f_j(\xi),$$

we have

$$\begin{aligned} \frac{\xi_0 \left(\mathcal{D}_s^{(\alpha, \beta)} F(\xi_0) \right)'}{\mathcal{D}_s^{(\alpha, \beta)} F(\xi_0)} &= \frac{1}{\vartheta} \left[\frac{\xi_0 \sum_{i=1}^{\vartheta} \left(\mathcal{D}_s^{(\alpha, \beta)} f_i(\xi_0) \right)'}{\frac{1}{\vartheta} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} f_j(\xi_0)} \right] \\ &= \frac{1}{\vartheta} \sum_{i=1}^{\vartheta} \mu(w_i) = \mu(w_0), \end{aligned}$$

for some $w_0 \in \mathcal{U}$, since μ is convex in $\mathcal{U}$. □

Putting $\mu(\xi) = \frac{1-\xi}{1+\xi}$ in Theorem 1 we obtain:

Corollary 1 If $f = \{f_1, f_2, \dots, f_{\vartheta}\} \in \mathfrak{R}_s^{(\alpha, \beta)}\left(\vartheta; \frac{1-\xi}{1+\xi}\right)$, then $\mathcal{D}_s^{(\alpha, \beta)} f_i(\xi)$ are close-to-convex univalent functions (see Kaplan [33]).

Theorem 2 Let $f = \{f_1, f_2, \dots, f_{\vartheta}\} \in \mathfrak{R}_s^{(\alpha, \beta)}(\vartheta; \mu)$. Define

$$F_i(\xi) = \frac{\varsigma+1}{\xi\varsigma} \int_0^{\xi} t^{\varsigma-1} f_i(t) dt \quad (\varsigma > 0; i = 1, 2, \dots, \vartheta).$$

If μ is bounded in $\mathcal{U}$ and $Re(\mu(\xi) + \varsigma) > 0$, then $F = \{F_1, F_2, \dots, F_{\vartheta}\} \in \mathfrak{R}_s^{(\alpha, \beta)}(\vartheta; \mu)$.

Proof. From the definition of $F_i(\xi)$, it follows that

$$\xi F_i'(\xi) + \varsigma F_i(\xi) = (\varsigma + 1)f_i(\xi),$$

and on taking convolution with $\mathfrak{D}_s^{(\alpha, \beta)}$ given by (9), we get

$$\xi [\mathcal{D}_s^{(\alpha, \beta)} F_i(\xi)]' + \varsigma \mathcal{D}_s^{(\alpha, \beta)} F_i(\xi) = (\varsigma + 1) \mathcal{D}_s^{(\alpha, \beta)} f_i(\xi), \quad i = 1, 2, \dots, \vartheta. \quad (10)$$

Let

$$p_i(\xi) = \frac{\vartheta \xi [\mathcal{D}_s^{(\alpha, \beta)} F_i(\xi)]'}{\sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} F_j(\xi)}, \quad (11)$$

where $\xi^{-1} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} F_j(\xi) \neq 0$. From (10), we have

$$\frac{p_i(\xi)}{\vartheta} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} F_j(\xi) + \varsigma \mathcal{D}_s^{(\alpha, \beta)} F_i(\xi) = (\varsigma + 1) \mathcal{D}_s^{(\alpha, \beta)} f_i(\xi). \quad (12)$$

Differentiating (12) with respect to ξ , we get

$$\frac{p_i'(\xi)}{\vartheta} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} F_j(\xi) + \frac{p_i(\xi)}{\vartheta} \sum_{j=1}^{\vartheta} [\mathcal{D}_s^{(\alpha, \beta)} F_j(\xi)]' + \varsigma [\mathcal{D}_s^{(\alpha, \beta)} F_i(\xi)]' = (\varsigma + 1) [\mathcal{D}_s^{(\alpha, \beta)} f_i(\xi)]'.$$

From (11), we have

$$\begin{aligned} & p_i'(\xi) \frac{\sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} F_j(\xi)}{\vartheta} + \frac{p_i(\xi)}{\vartheta} \frac{\sum_{i=1}^{\vartheta} p_i(\xi) \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} F_j(\xi)}{\vartheta \xi} + \varsigma \frac{p_i(\xi) \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} F_j(\xi)}{\vartheta \xi} \\ & = (\varsigma + 1) [\mathcal{D}_s^{(\alpha, \beta)} f_i(\xi)]'. \end{aligned}$$

Hence

$$p_i'(\xi) + \frac{p_i(\xi)}{\vartheta \xi} \sum_{i=1}^{\vartheta} p_i(\xi) + \varsigma \frac{p_i(\xi)}{\xi} = \frac{(\varsigma + 1) [\mathcal{D}_s^{(\alpha, \beta)} f_i(\xi)]'}{\frac{1}{\vartheta} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} F_j(\xi)}.$$

Then

$$\begin{aligned} \frac{\xi p'_i(\xi)}{\frac{1}{\vartheta} \sum_{i=1}^{\vartheta} p_i(\xi) + \varsigma} + p_i(\xi) &= \frac{(\varsigma + 1) \xi [\mathcal{D}_s^{(\alpha, \beta)} f_i(\xi)]'}{\frac{1}{\vartheta} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} F_j(\xi)} \cdot \frac{1}{\frac{1}{\vartheta} \sum_{i=1}^{\vartheta} p_i(\xi) + \varsigma} \\ &= \frac{(\varsigma + 1) \xi [\mathcal{D}_s^{(\alpha, \beta)} f_i(\xi)]'}{\frac{1}{\vartheta} \left\{ \frac{1}{\vartheta} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} F_j(\xi) \cdot \sum_{i=1}^{\vartheta} p_i(\xi) + \varsigma \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} F_j(\xi) \right\}}. \end{aligned}$$

From (12), we have

$$\frac{\xi p'_i(\xi)}{\frac{1}{\vartheta} \sum_{i=1}^{\vartheta} p_i(\xi) + \varsigma} + p_i(\xi) = \frac{(\varsigma + 1) \xi [\mathcal{D}_s^{(\alpha, \beta)} f_i(\xi)]'}{\frac{1}{\vartheta} (\varsigma + 1) \sum_{i=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} f_i(\xi)} \prec \mu(\xi), \quad (13)$$

since $f = \{f_1, f_2, \dots, f_{\vartheta}\} \in \mathfrak{R}_s^{(\alpha, \beta)}(\vartheta; \mu)$. Now we can write for any $\xi_0 \in \mathcal{U}$,

$$\frac{\frac{1}{\vartheta} \xi_0 p'_i(\xi_0)}{\frac{1}{\vartheta} \sum_{j=1}^{\vartheta} p_j(\xi_0) + \varsigma} + \frac{1}{\vartheta} p_i(\xi_0) = \frac{1}{\vartheta} \mu(w_i),$$

for some $w_i \in \mathcal{U}$. This is true for $i = 1, 2, \dots, \vartheta$. Since μ is convex, there exists a $w_0 \in \mathcal{U}$ such that

$$\frac{\xi_0 Q'(\xi_0)}{Q(\xi_0) + \varsigma} + Q(\xi_0) = \mu(w_0),$$

where $Q(\xi) = \frac{1}{\vartheta} \sum_{i=1}^{\vartheta} p_i(\xi)$. Hence

$$\frac{\xi Q'(\xi)}{Q(\xi) + \varsigma} + Q(\xi) \prec \mu(\xi). \quad (14)$$

Since $Re\{\mu\}$ is bounded and $Re(\mu(\xi) + \varsigma) > 0$. By Lemma 1 with $Q(\xi)$ in (14) we obtain $Q(\xi) \prec \mu(\xi)$ ($\xi \in \mathcal{U}$). From (13), we have

$$\frac{\xi p'_i(\xi)}{Q(\xi) + \varsigma} + p_i(\xi) \prec \mu(\xi), \quad (15)$$

where $Q(\xi) \prec \mu(\xi)$. By Lemma 2 with $\mu(\xi)$ in (15) we obtain $p_i(\xi) \prec \mu(\xi)$ ($\xi \in \mathcal{U}$), $i = 1, 2, \dots, \vartheta$, that is

$$\frac{\xi[\mathcal{D}_s^{(\alpha, \beta)} F_i(\xi)]'}{\frac{1}{\vartheta} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} F_j(\xi)} \prec \mu(\xi).$$

Now

$$F_i(\xi) = \frac{\varsigma+1}{\xi^\varsigma} \int_0^\xi t^{\varsigma-1} f_i(t) dt, \quad (\varsigma > 0).$$

For every i , $1 \leq i \leq \vartheta$,

$$\mathcal{D}_s^{(\alpha, \beta)} F_i(\xi) = \frac{\varsigma+1}{\xi^\varsigma} \int_0^\xi t^{\varsigma-1} \mathcal{D}_s^{(\alpha, \beta)} f_i(t) dt,$$

and hence

$$\begin{aligned} \sum_{i=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} F_i(\xi) &= \frac{\varsigma+1}{\xi^\varsigma} \int_0^\xi t^{\varsigma-1} \sum_{i=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} f_i(t) dt \\ &= \frac{\varsigma+1}{\xi^\varsigma} \int_0^\xi t^\varsigma g(t) dt, \end{aligned}$$

where $g(t) = t^{-1} \sum_{i=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} f_i(t) \neq 0$, for $\xi \in \mathcal{U}$. Now define

$$\Omega(\xi) = \sum_{\kappa=1}^{\infty} \frac{\varsigma+1}{\varsigma+\kappa} \xi^{\kappa-1}, \quad \operatorname{Re}(\varsigma) > 0.$$

Then an easy calculations show that

$$\xi^{-1} \sum_{i=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} F_i(\xi) = (\Omega * g)(\xi) \neq 0.$$

Thus $F = \{F_1, F_2, \dots, F_\vartheta\} \in \mathfrak{A}_s^{(\alpha, \beta)}(\vartheta; \mu)$. □

Remark 2 (i) Let $\vartheta = s = 1$, and $\alpha = \beta = 0$ in Theorem 2, we get the result obtained by Padmanabhan and Parvatham ([30], Theorem 2 with $a = 1$);

(ii) Let $\vartheta = s = \varsigma = 1$, $\alpha = \beta = 0$ and $\mu(\xi) = \frac{1-\xi}{1+\xi}$, in Theorem 2, we get the result obtained by Libera ([33], Theorem 1);

(iii) Let $\vartheta = s = 1$, $\alpha = \beta = 0$ and $\mu(\xi) = \frac{1+A\xi}{1+B\xi}$, $(-1 \leq B < A \leq 1)$, in Theorem 2, we get the result obtained by Goel and Mehrok ([34], Lemma 3).

Theorem 3 If $f = \{f_1, f_2, \dots, f_\vartheta\} \in \mathfrak{R}_s^{(\alpha+\frac{1}{s}, \beta-\frac{1}{s})}(\vartheta; \mu)$, and $Re\{\mu\}$ is bounded in $\mathcal{U}$, then $f = \{f_1, f_2, \dots, f_\vartheta\} \in \mathfrak{R}_s^{(\alpha, \beta)}(\vartheta; \mu)$ holds for $Re[\mu(\xi) + (s(1-\alpha) - 1)] > 0$ in $\mathcal{U}$.

Proof. Let

$$p_i(\xi) = \frac{\vartheta \xi [\mathcal{D}_s^{(\alpha, \beta)} f_i(\xi)]'}{\sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} f_j(\xi)} \quad (\xi \in \mathcal{U}; i = 1, 2, \dots, \vartheta), \quad (16)$$

where $\xi^{-1} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} f_j(\xi) \neq 0$. From (7) and (16), we have

$$\frac{1}{\vartheta} p_i(\xi) \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} f_j(\xi) = s(1-\alpha) \mathcal{D}_s^{(\alpha+\frac{1}{s}, \beta-\frac{1}{s})} f_i(\xi) - [s(1-\alpha) - 1] \mathcal{D}_s^{(\alpha, \beta)} f_i(\xi). \quad (17)$$

Differentiating (17) with respect to ξ , we get

$$\begin{aligned} \frac{\xi}{\vartheta} p_i'(\xi) \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} f_j(\xi) + \frac{\xi}{\vartheta} p_i(\xi) \sum_{j=1}^{\vartheta} [\mathcal{D}_s^{(\alpha, \beta)} f_j(\xi)]' &= s(1-\alpha) \xi [\mathcal{D}_s^{(\alpha+\frac{1}{s}, \beta-\frac{1}{s})} f_i(\xi)]' \\ &\quad - [s(1-\alpha) - 1] \xi [\mathcal{D}_s^{(\alpha, \beta)} f_i(\xi)]'. \end{aligned}$$

Using (16), we obtain

$$\begin{aligned} \frac{\xi}{\vartheta} p_i'(\xi) \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} f_j(\xi) + p_i(\xi) \left[\frac{\xi}{\vartheta} \sum_{j=1}^{\vartheta} [\mathcal{D}_s^{(\alpha, \beta)} f_j(\xi)]' + \frac{[s(1-\alpha) - 1]}{\vartheta} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} f_j(\xi) \right] \\ = s(1-\alpha) \xi [\mathcal{D}_s^{(\alpha+\frac{1}{s}, \beta-\frac{1}{s})} f_i(\xi)]'. \end{aligned}$$

Then

$$\frac{\frac{\xi}{\vartheta} p'_i(\xi) \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} f_j(\xi)}{\frac{\xi}{\vartheta} \sum_{j=1}^{\vartheta} [\mathcal{D}_s^{(\alpha, \beta)} f_j(\xi)]' + \frac{[s(1-\alpha)-1]}{\vartheta} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} f_j(\xi)} + p_i(\xi)$$

$$= \frac{s(1-\alpha)\xi [\mathcal{D}_s^{(\alpha+\frac{1}{s}, \beta-\frac{1}{s})} f_i(\xi)]'}{\frac{\xi}{\vartheta} \sum_{j=1}^{\vartheta} [\mathcal{D}_s^{(\alpha, \beta)} f_j(\xi)]' + \frac{[s(1-\alpha)-1]}{\vartheta} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} f_j(\xi)}.$$

Using (7), we have

$$\frac{\frac{\xi}{\vartheta} p'_i(\xi) \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} f_j(\xi)}{\frac{\xi}{\vartheta} \sum_{j=1}^{\vartheta} [\mathcal{D}_s^{(\alpha, \beta)} f_j(\xi)]' + \frac{[s(1-\alpha)-1]}{\vartheta} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} f_j(\xi)} + p_i(\xi) = \frac{\xi [\mathcal{D}_s^{(\alpha+\frac{1}{s}, \beta-\frac{1}{s})} f_i(\xi)]'}{\frac{1}{\vartheta} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha+\frac{1}{s}, \beta-\frac{1}{s})} f_j(\xi)}. \quad (18)$$

Using (16) in (18), we have

$$\frac{\xi p'_i(\xi)}{\frac{1}{\vartheta} \sum_{j=1}^{\vartheta} p_j(\xi) + [s(1-\alpha)-1]} + p_i(\xi) = \frac{\xi [\mathcal{D}_s^{(\alpha+\frac{1}{s}, \beta-\frac{1}{s})} f_i(\xi)]'}{\frac{1}{\vartheta} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha+\frac{1}{s}, \beta-\frac{1}{s})} f_j(\xi)}.$$

Since $f = \{f_1, f_2, \dots, f_{\vartheta}\} \in \mathfrak{R}_s^{(\alpha+\frac{1}{s}, \beta-\frac{1}{s})}(\vartheta; \mu)$, then we have

$$\frac{\xi p'_i(\xi)}{\frac{1}{\vartheta} \sum_{j=1}^{\vartheta} p_j(\xi) + [s(1-\alpha)-1]} + p_i(\xi) = \frac{\xi [\mathcal{D}_s^{(\alpha+\frac{1}{s}, \beta-\frac{1}{s})} f_i(\xi)]'}{\frac{1}{\vartheta} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha+\frac{1}{s}, \beta-\frac{1}{s})} f_j(\xi)} \prec \mu(\xi), i = 1, 2, \dots, \vartheta. \quad (19)$$

Therefore for any $\xi_0 \in \mathcal{U}$, we have

$$\frac{\frac{1}{\vartheta} \xi_0 p'_i(\xi_0)}{\frac{1}{\vartheta} \sum_{j=1}^{\vartheta} p_j(\xi_0) + [s(1-\alpha)-1]} + \frac{1}{\vartheta} p_i(\xi_0) = \frac{1}{\vartheta} \mu(w_i)$$

for some $w_0 \in \mathcal{U}$. Since μ is convex, there exists a $w_i \in \mathcal{U}$, such that

$$\frac{\frac{\xi_0}{\vartheta} \sum_{i=1}^n p'_i(\xi_0)}{\frac{1}{\vartheta} \sum_{j=1}^{\vartheta} p_j(\xi_0) + [s(1-\alpha) - 1]} + \frac{1}{\vartheta} \sum_{j=1}^{\vartheta} p_j(\xi_0) = \frac{1}{\vartheta} \sum_{i=1}^{\vartheta} \mu(w_i) = \mu(w_0).$$

Setting $Q(\xi) = \frac{1}{\vartheta} \sum_{i=1}^{\vartheta} p_i(\xi)$, we have

$$\frac{\xi Q'(\xi)}{Q(\xi) + [s(1-\alpha) - 1]} + Q(\xi) \prec \mu(\xi), \quad (20)$$

by Lemma 1 with $Q(\xi)$ in (20) we obtain $Q(\xi) \prec \mu(\xi)$. (19) gives us

$$\frac{\xi p'_i(\xi)}{Q(\xi) + [s(1-\alpha) - 1]} + p_i(\xi) \prec \mu(\xi), \quad (21)$$

by Lemma 2 with $\mu(\xi)$ in (21) we obtain $p_i(\xi) \prec \mu(\xi)$, which implies that $f = \{f_1, f_2, \dots, f_{\vartheta}\} \in \mathfrak{R}_s^{(\alpha, \beta)}(\vartheta; \mu)$. $\square$

Remark 3 Let $\vartheta = s = 1$, and $\alpha = \beta = 0$, in Theorem 3, we get the result obtained by Padmanabhan and Parvathem ([30], Theorem 1 with $a = 1$);

2.2 The class $\mathfrak{C}_s^{(\alpha, \beta)}(\vartheta; \mu)$

Definition 2 Let $\mathfrak{C}_s^{(\alpha, \beta)}(\vartheta; \mu)$ denote the class of functions $f \in \mathcal{A}$ which satisfies

$$\frac{\xi [\mathcal{D}_s^{(\alpha, \beta)} f(\xi)]'}{\frac{1}{\vartheta} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} g_j(\xi)} \prec \mu(\xi) \quad (\xi \in \mathcal{U}),$$

where $\xi^{-1} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} g_j(\xi) \neq 0$ and $g = \{g_1, g_2, \dots, g_{\vartheta}\} \in \mathfrak{R}_s^{(\alpha, \beta)}(\vartheta; \mu)$, μ is convex univalent in $\mathcal{U}$ with $\mu(0) = 1$.

Remark 4 (i) Let $\vartheta = s = 1$, $\alpha = \beta = 0$ and $\mu(\xi) = \frac{1-\xi}{1+\xi}$, then $\mathfrak{C}_1^{(0, 0)}\left(1; \frac{1-\xi}{1+\xi}\right) = \mathfrak{C}$, where $\mathfrak{C}$ is the class of close-to-convex functions (see [35]);

(ii) Let $\vartheta = s = 1$, $\alpha = \beta = 0$ and $\mu(\xi) = \frac{1+A\xi}{1+B\xi}$, $(-1 \leq B < A \leq 1)$, then $\mathfrak{C}_1^{(0, 0)}\left(1; \frac{1+A\xi}{1+B\xi}\right) = \mathfrak{C}(A, B)$, where $\mathfrak{C}(A, B)$ is the generalized class of close-to-convex functions (see [36, 37]).

Theorem 4 Let $f \in \mathfrak{C}_s^{(\alpha, \beta)}(\vartheta; \mu)$. If $Re(\mu)$ is bounded in $\mathcal{U}$ and $Re(\mu(\xi) + \tau) > 0$, then

$$F(\xi) = \frac{\tau+1}{\xi^\tau} \int_0^\xi t^{\tau-1} f(t) dt \quad (\xi \in \mathcal{U}; \tau > 0),$$

also belongs to $\mathfrak{C}_s^{(\alpha, \beta)}(\vartheta; \mu)$.

Proof. Since $f \in \mathfrak{C}_s^{(\alpha, \beta)}(\vartheta; \mu)$, then there exists $g = \{g_1, g_2, \dots, g_\vartheta\} \in \mathfrak{R}_s^{(\alpha, \beta)}(\vartheta; \mu)$, such that

$$\frac{\xi [\mathcal{D}_s^{(\alpha, \beta)} f(\xi)]'}{\frac{1}{\vartheta} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} g_j(\xi)} \prec \mu(\xi) \quad (\xi \in \mathcal{U}),$$

where $\xi^{-1} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} g_j(\xi) \neq 0$. Let

$$G_i(\xi) = \frac{\tau+1}{\xi^\tau} \int_0^\xi t^{\tau-1} g_i(t) dt \quad (Re\tau > 0).$$

Then by Theorem 2, we have $G = \{G_1, G_2, \dots, G_\vartheta\} \in \mathfrak{R}_s^{(\alpha, \beta)}(\vartheta; \mu)$. Also let

$$p(\xi) = \frac{\xi [\mathcal{D}_s^{(\alpha, \beta)} F(\xi)]'}{\frac{1}{\vartheta} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} G_j(\xi)} \quad (\xi \in \mathcal{U}). \quad (22)$$

Now, from the definitions of G_i and F , we have

$$\xi [\mathcal{D}_s^{(\alpha, \beta)} G_i(\xi)]' + \tau \mathcal{D}_s^{(\alpha, \beta)} G_i(\xi) = (\tau+1) \mathcal{D}_s^{(\alpha, \beta)} g_i(\xi), \quad (23)$$

and

$$\xi [\mathcal{D}_s^{(\alpha, \beta)} F(\xi)]' + \tau \mathcal{D}_s^{(\alpha, \beta)} F(\xi) = (\tau+1) \mathcal{D}_s^{(\alpha, \beta)} f(\xi). \quad (24)$$

From (22) to (24), we have

$$\frac{1}{\vartheta} p(\xi) \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} G_j(\xi) + \tau \mathcal{D}_s^{(\alpha, \beta)} F(\xi) = (\tau+1) \mathcal{D}_s^{(\alpha, \beta)} f(\xi). \quad (25)$$

Differentiating (25) with respect to ξ , and multiplying the resulting equation by ξ , we have

$$\frac{\xi}{\vartheta} p'(\xi) \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} G_j(\xi) + \frac{\xi}{\vartheta} p(\xi) \sum_{j=1}^{\vartheta} [\mathcal{D}_s^{(\alpha, \beta)} G_j(\xi)]' + \tau \xi [\mathcal{D}_s^{(\alpha, \beta)} F(\xi)]' = (\tau+1) \xi [\mathcal{D}_s^{(\alpha, \beta)} f(\xi)]'. \quad (26)$$

From (22) into (26), we have

$$\frac{\xi}{\vartheta} p'(\xi) \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} G_j(\xi) + \frac{\xi}{\vartheta} p(\xi) \sum_{j=1}^{\vartheta} [\mathcal{D}_s^{(\alpha, \beta)} G_j(\xi)]' + \tau \frac{p(\xi)}{\vartheta} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} G_j(\xi) = (\tau + 1) \xi [\mathcal{D}_s^{(\alpha, \beta)} f(\xi)]'.$$

Hence, we get

$$\begin{aligned} & \frac{\frac{\xi}{\vartheta} p'(\xi) \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} G_j(\xi)}{\frac{\xi}{\vartheta} \sum_{j=1}^{\vartheta} [\mathcal{D}_s^{(\alpha, \beta)} G_j(\xi)]' + \frac{\tau}{\vartheta} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} G_j(\xi)} + p(\xi) \\ &= \frac{(\tau + 1) \xi [\mathcal{D}_s^{(\alpha, \beta)} f(\xi)]'}{\frac{\xi}{\vartheta} \sum_{j=1}^{\vartheta} [\mathcal{D}_s^{(\alpha, \beta)} G_j(\xi)]' + \frac{\tau}{\vartheta} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} G_j(\xi)} = \frac{\xi [\mathcal{D}_s^{(\alpha, \beta)} f(\xi)]'}{\frac{1}{\vartheta} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} g_j(\xi)} \quad (\text{by using (23)}). \end{aligned}$$

From the above, we have

$$\frac{\xi p'(\xi)}{\frac{1}{\vartheta} \sum_{j=1}^{\vartheta} Q_j(\xi) + \tau} + p(\xi) = \frac{\xi [\mathcal{D}_s^{(\alpha, \beta)} f(\xi)]'}{\frac{1}{\vartheta} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} g_j(\xi)} \prec \mu(\xi),$$

where $Q_j(\xi) = \frac{\xi [\mathcal{D}_s^{(\alpha, \beta)} G_j(\xi)]'}{\frac{1}{\vartheta} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} G_j(\xi)}$. Now $Q_j(\xi) \prec \mu(\xi)$, $j = 1, 2, \dots, \vartheta$, since $G = \{G_1, G_2, \dots, G_{\vartheta}\} \in \mathfrak{R}_s^{(\alpha, \beta)}(\vartheta; \mu)$

and μ is a convex univalent. Since $Re(\mu(\xi) + \tau) > 0$, Lemma 2 applied indicates that $p(\xi) \prec \mu(\xi)$, hence $F \in \mathfrak{C}_s^{(\alpha, \beta)}(\vartheta; \mu)$. $\square$

Remark 5 (i) Let $\vartheta = s = 1$, and $\alpha = \beta = 0$ in Theorem 4, we get the result obtained by Padmanabhan and Parvathem ([30], Theorem 4 with $a = 1$);

(ii) Let $\vartheta = s = 1$, $\alpha = \beta = 0$ and $\mu(\xi) = \frac{1 - \xi}{1 + \xi}$, in Theorem 2, we get the result obtained by Libera ([33], Theorem 1).

Theorem 5 If $f \in \mathfrak{C}_s^{(\alpha + \frac{1}{s}, \beta - \frac{1}{s})}(\vartheta; \mu)$ and μ is bounded in $\mathcal{U}$, then $f \in \mathfrak{C}_s^{(\alpha, \beta)}(\vartheta; \mu)$ holds for $Re[\mu(\xi) + (s(1 - \alpha) - 1)] > 0$ in $\mathcal{U}$.

Proof. The proof of the theorem is similar to the proof of Theorem 3. $\square$

Remark 6 Let $\vartheta = s = 1$, and $\alpha = \beta = 0$ in Theorem 5, we get the result obtained by Padmanabhan and Parvathem ([30], Theorem 3 with $a = 1$);

2.3 The class $\mathfrak{P}_s^{(\alpha, \beta)}(\vartheta; \delta, \mu)$

Definition 3 Let $\mathfrak{P}_s^{(\alpha, \beta)}(\vartheta; \delta, \mu)$, $\delta \geq 0$, denote the class of function $f \in \mathcal{A}$ satisfies

$$\mathfrak{J}(\delta; f; g_1, g_2, \dots, g_{\vartheta})(\xi) = \left\{ \delta \frac{\xi \left[\mathcal{D}_s^{(\alpha+\frac{1}{s}, \beta-\frac{1}{s})} f(\xi) \right]'}{\frac{1}{\vartheta} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha+\frac{1}{s}, \beta-\frac{1}{s})} g_j(\xi)} + (1-\delta) \frac{\xi \left[\mathcal{D}_s^{(\alpha, \beta)} f(\xi) \right]'}{\frac{1}{\vartheta} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} g_j(\xi)} \right\} \prec \mu(\xi) \quad (\xi \in \mathcal{U}),$$

where $g = \{g_1, g_2, \dots, g_{\vartheta}\} \in \mathfrak{R}_s^{(\alpha, \beta)}(\vartheta; \mu)$, $\xi^{-1} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha+\frac{1}{s}, \beta-\frac{1}{s})} g_j(\xi) \neq 0$ in $\mathcal{U}$, μ is convex univalent in $\mathcal{U}$ with $\mu(0) = 1$.

Remark 7 We note that $\mathcal{P}_s^{(\alpha, \beta)}(\vartheta; 0, \mu) = \mathfrak{C}_s^{(\alpha, \beta)}(\vartheta; \mu)$.

Theorem 6 If $f \in \mathcal{P}_s^{(\alpha, \beta)}(\vartheta; \delta, \mu)$ and $Re\{\mu\}$ is bounded in $\mathcal{U}$, then $f \in \mathcal{P}_s^{(\alpha, \beta)}(\vartheta; 0, \mu) = \mathfrak{C}_s^{(\alpha, \beta)}(\vartheta; \mu)$ hold for $Re[\mu(\xi) + (s(1-\alpha)-1)] > 0$.

Proof. For $\delta = 0$, the theorem is unimportant, we can presume that $\delta \neq 0$. Let

$$p(\xi) = \frac{\xi \left[\mathcal{D}_s^{(\alpha, \beta)} f(\xi) \right]'}{\frac{1}{\vartheta} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} g_j(\xi)} \quad (\xi \in \mathcal{U}),$$

where $\xi^{-1} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} g_j(\xi) \neq 0$. Then an easy calculations shows that

$$\frac{\xi p'(\xi)}{\frac{1}{\vartheta} \sum_{j=1}^{\vartheta} q_j(\xi) + [s(1-\alpha)-1]} + p(\xi) = \frac{\xi \left[\mathcal{D}_s^{(\alpha+\frac{1}{s}, \beta-\frac{1}{s})} f(\xi) \right]'}{\frac{1}{\vartheta} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha+\frac{1}{s}, \beta-\frac{1}{s})} g_j(\xi)},$$

where $q_j(\xi) = \frac{\xi \left[\mathcal{D}_s^{(\alpha, \beta)} g_j(\xi) \right]'}{\frac{1}{\vartheta} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} g_j(\xi)}$. Also $\frac{1}{\vartheta} \sum_{j=1}^{\vartheta} q_j(\xi) \prec \mu(\xi)$. Since $f \in \mathcal{P}_s^{(\alpha, \beta)}(\vartheta; \delta, \mu)$, we have

$$\mathfrak{J}(\delta; f; g_1, g_2, \dots, g_{\vartheta})(\xi) = \frac{\delta \xi p'(\xi)}{\frac{1}{\vartheta} \sum_{j=1}^{\vartheta} q_j(\xi) + [s(1-\alpha)-1]} + p(\xi) \prec \mu(\xi). \quad (27)$$

By Lemma 2 with $\mu(\xi)$ in (27) we obtain $p(\xi) \prec \mu(\xi)$ which implies $f \in \mathfrak{C}_s^{(\alpha, \beta)}(\vartheta; \mu)$. $\square$

Remark 8 (i) Let $\vartheta = s = 1$, and $\alpha = \beta = 0$ in Theorem 6, we get the result obtained by Padmanabhan and Parvatham ([30], Theorem 5 with $a = 1$);

(ii) Let $\vartheta = s = 1$, $\alpha = \beta = 0$ and $\mu(\xi) = \frac{1-\xi}{1+\xi}$, in Theorem 6 we get the result obtained by Chichra ([38], Theorem 1);

(iii) Let $\vartheta = s = 1$, $\alpha = \beta = 0$ and $\mu(\xi) = \frac{1+(2\rho-1)\xi}{1+\xi}$ ($0 \leq \rho < 1$), in Theorem 6 we get the result obtained by Zmorovich and Pokhilevich ([39], Theorem 1).

Theorem 7 For $\delta > \lambda \geq 0$ and $Re\mu(\xi)$ is bounded in $\mathcal{U}$, then $\wp_s^{(\alpha, \beta)}(\vartheta; \delta, \mu) \subset \wp_s^{(\alpha, \beta)}(\vartheta; \lambda, \mu)$.

Proof. The case $\lambda = 0$ was treated in the previous theorem. Hence we assume that $\lambda \neq 0$. Suppose that $f \in \wp_s^{(\alpha, \beta)}(\vartheta; \delta, \mu)$. Then

$$\mathfrak{J}(\delta; f; g_1, g_2, \dots, g_\vartheta)(\xi) \prec \mu(\xi). \quad (28)$$

Let ξ_1 be any arbitrary point in $\mathcal{U}$. Then

$$\mathfrak{J}(\delta; f; g_1, g_2, \dots, g_\vartheta)(\xi_1) \prec \mu(\mathcal{U}).$$

From Theorem 6, we have

$$\frac{\xi \left[\mathcal{D}_s^{(\alpha, \beta)} f(\xi) \right]'}{\frac{1}{\vartheta} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} g_j(\xi)} \prec \mu(\xi). \quad (29)$$

Now

$$\mathfrak{J}(\lambda; f; g_1, g_2, \dots, g_\vartheta)(\xi) = \left(1 - \frac{\lambda}{\delta}\right) \frac{\xi \left[\mathcal{D}_s^{(\alpha, \beta)} f(\xi) \right]'}{\frac{1}{\vartheta} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} g_j(\xi)} + \frac{\lambda}{\delta} \mathfrak{J}(\delta; f; g_1, g_2, \dots, g_\vartheta)(\xi).$$

From (28) and (29) it follows that

$$\frac{\xi_1 \left[\mathcal{D}_s^{(\alpha, \beta)} f(\xi_1) \right]'}{\frac{1}{\vartheta} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} g_j(\xi_1)} \prec \mu(\mathcal{U})$$

and

$$\delta \frac{\xi_1 \left[\mathcal{D}_s^{(\alpha + \frac{1}{s}, \beta - \frac{1}{s})} f(\xi_1) \right]'}{\frac{1}{\vartheta} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha + \frac{1}{s}, \beta - \frac{1}{s})} g_j(\xi_1)} + (1 - \delta) \frac{\xi_1 \left[\mathcal{D}_s^{(\alpha, \beta)} f(\xi_1) \right]'}{\frac{1}{\vartheta} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} g_j(\xi_1)} \prec \mu(\mathcal{U}).$$

Now $\mu(\mathcal{U})$ is convex and $\frac{\lambda}{\delta} < 1$, hence we have $\mathfrak{J}(\lambda; f; g_1, g_2, \dots, g_\vartheta)(\xi_1) \prec \mu(\mathcal{U})$, showing that $f \in \wp_s^{(\alpha, \beta)}(\vartheta; \lambda, \mu)$. $\square$

Remark 9 (i) Let $\vartheta = s = 1$, and $\alpha = \beta = 0$ in Theorem 7, Padmanabhan and Parvatham ([30], Theorem 6 with $a = 1$);

(ii) Let $\vartheta = s = 1$, $\alpha = \beta = 0$ and $\mu(\xi) = \frac{1-\xi}{1+\xi}$, in Theorem 7 we obtain the result obtained by Chichra ([38], Theorem 2);

(iii) Let $\vartheta = s = 1$, $\alpha = \beta = 0$ and $\mu(\xi) = \frac{1+(2\rho-1)\xi}{1+\xi}$ ($0 \leq \rho < 1$), in Theorem 7 we obtain the result obtained by Zmorovich and Pokhilevich ([39], Theorem 1).

2.4 The class $\mathfrak{R}_s^{(\alpha, \beta)}(\vartheta; \delta, \mu)$

Definition 4 Let $\mathfrak{R}_s^{(\alpha, \beta)}(\vartheta; \delta, \mu)$, $\delta \geq 0$, denote the class of function $f \in \mathcal{A}$ satisfies

$$\mathfrak{J}(\delta; f; f_1, f_2, \dots, f_\vartheta)(\xi) = \left\{ \delta \frac{\xi \left[\mathcal{D}_s^{(\alpha+\frac{1}{s}, \beta-\frac{1}{s})} f_i(\xi) \right]'}{\frac{1}{\vartheta} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha+\frac{1}{s}, \beta-\frac{1}{s})} f_j(\xi)} + (1-\delta) \frac{\xi \left[\mathcal{D}_s^{(\alpha, \beta)} f_i(\xi) \right]'}{\frac{1}{\vartheta} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} f_j(\xi)} \right\} \prec \mu(\xi) \quad (\xi \in \mathcal{U}),$$

where $f = \{f_1, f_2, \dots, f_\vartheta\} \in \mathfrak{R}_s^{(\alpha, \beta)}(\vartheta; \mu)$ and $\xi^{-1} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha+\frac{1}{s}, \beta-\frac{1}{s})} f_j(\xi) \neq 0$ in $\mathcal{U}$, μ is convex univalent in $\mathcal{U}$ with $\mu(0) = 1$.

We note that $\mathfrak{R}_s^{(\alpha, \beta)}(\vartheta; 0; \mu) = \mathfrak{R}_s^{(\alpha, \beta)}(\vartheta; \mu)$.

Theorem 8 If $f \in \mathfrak{R}_s^{(\alpha, \beta)}(\vartheta; \delta; \mu)$ and $Re\{\mu\}$ is bounded in $\mathcal{U}$, then $f \in \mathfrak{R}_s^{(\alpha, \beta)}(\vartheta; 0; \mu) = \mathfrak{R}_s^{(\alpha, \beta)}(\vartheta; \mu)$ hold for $Re[\mu(\xi) + (s(1-\alpha)-1)] > 0$.

Proof. For $\delta = 0$, the theorem is unimportant, we can presume that $\delta \neq 0$. Let

$$p(\xi) = \frac{\xi \left[\mathcal{D}_s^{(\alpha, \beta)} f_i(\xi) \right]'}{\frac{1}{\vartheta} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} f_j(\xi)} \quad (\xi \in \mathcal{U}),$$

where $\xi^{-1} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} f_j(\xi) \neq 0$. Then an easy calculations shows that

$$\frac{\xi p'(\xi)}{\frac{1}{\vartheta} \sum_{j=1}^{\vartheta} q_j(\xi) + [s(1-\alpha)-1]} + p(\xi) = \frac{\xi \left[\mathcal{D}_s^{(\alpha+\frac{1}{s}, \beta-\frac{1}{s})} f_i(\xi) \right]'}{\frac{1}{\vartheta} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha+\frac{1}{s}, \beta-\frac{1}{s})} f_j(\xi)},$$

where $q_j(\xi) = \frac{\xi \left[\mathcal{D}_s^{(\alpha, \beta)} f_j(\xi) \right]'}{\frac{1}{\vartheta} \sum_{j=1}^{\vartheta} \mathcal{D}_s^{(\alpha, \beta)} f_j(\xi)}$. Also $\frac{1}{\vartheta} \sum_{j=1}^{\vartheta} q_j(\xi) \prec \mu(\xi)$. Since $f(\xi) \in \mathfrak{R}_s^{(\alpha, \beta)}(\vartheta; \delta; \mu)$, we have

$$J(\delta; f; f_1, f_2, \dots, f_{\vartheta})(\xi) = \frac{\delta \xi p'(\xi)}{\frac{1}{\vartheta} \sum_{j=1}^{\vartheta} q_j(\xi) + [s(1-\alpha) - 1]} + p(\xi) \prec \mu(\xi). \quad (30)$$

By Lemma 2 with $\mu(\xi)$ in (30) we obtain $p(\xi) \prec \mu(\xi)$ which implies $f \in \mathfrak{R}_s^{(\alpha, \beta)}(\vartheta; \mu)$. $\square$

Theorem 9 For $\delta > \lambda \geq 0$ and $Re\{\mu\}$ is bounded in $\mathcal{U}$, then $\mathfrak{R}_s^{(\alpha, \beta)}(\vartheta; \delta; \mu) \subset \mathfrak{R}_s^{(\alpha, \beta)}(\vartheta; \lambda; \mu)$.

Proof. The proof of the theorem is similar to the proof of Theorem 7. $\square$

Remark 10 (i) Let $\vartheta = s = 1$, and $\alpha = \beta = 0$ in Theorem 8, Padmanabhan and Parvathem [30], Theorem 7 with $a = 1$;

(ii) Let $\vartheta = s = 1$, $\alpha = \beta = 0$ and $\mu(\xi) = \frac{1-\xi}{1+\xi}$, in Theorem 8 we get the result obtained by Al-Amiri [40], Theorem 1;

(iii) Let $\vartheta = s = 1$, $\alpha = \beta = 0$ and $\mu(\xi) = \frac{1+(2\rho-1)\xi}{1+\xi}$ ($0 \leq \rho < 1$), in Theorem 8 we obtain the result obtained by Miller et al. ([41], Theorem 1).

3. Conclusion

As more applications and theoretical insights are discovered, the Zeta-Riemann fractional differential operator is poised to become an essential instrument in the modern mathematical toolkit. To investigate some classes of univalent functions, the theory of differential subordination is a result of the new findings in this paper. The idea for the study needed, Zeta- Riemann Fractional differential operator $\mathcal{D}_s^{(\alpha, \beta)} f(\xi)$, and the rationale behind the topic's investigation are all contained in Section 1's Introduction. This operator is applied in this investigation to define and investigate certain specific classes of univalent functions using the theory of differential subordination. The main findings are found in Sections 2, 3, 4, 6, where classes $\mathfrak{R}_s^{(\alpha, \beta)}(\vartheta; \mu)$, given in Definition 1, $\mathfrak{C}_s^{(\alpha, \beta)}(\vartheta; \mu)$, described in Definition 2, $\mathfrak{P}_s^{(\alpha, \beta)}(\vartheta; \delta, \mu)$ seen in Definition 3 and $\mathfrak{R}_s^{(\alpha, \beta)}(\vartheta; \delta, \mu)$, presented in Definition 4 are investigated, respectively. Also, we proved combinations of functions belonging to those classes and inclusion relations.

Acknowledgments

This research has been funded by Scientific Research Deanship at University of Hail-Saudi Arabia through project number RG-25006.

Data availability statement

The data used to support the findings of this study are included within the article.

Conflict of interest

The authors declare no competing financial interest.

References

- [1] Duren PL. *Univalent Functions*. New York: Springer; 1983.
- [2] Goodman AW. *Univalent Functions*. USA: Mancorp Pub; 1983.
- [3] Littlewood JE. *Lectures on the Theory of Functions*. London: Oxford University; 1944.
- [4] Rogosinski W. On coefficients of subordinate functions. *Proceedings of the London Mathematical Society*. 1945; s2-48(1): 48-82. Available from: [https://doi.org/10.1016/S0960-0779\(02\)00653-7](https://doi.org/10.1016/S0960-0779(02)00653-7).
- [5] Srivastava HM, Owa S. *Current Topics in Analytic Functions Theory*. Singapore: World Scientific; 1992. Available from: <https://doi.org/10.1142/1628>.
- [6] Miller SS, Mocanu PT. Second order differential inequalities in the complex plane. *Journal of Mathematical Analysis and Applications*. 1978; 65(2): 298-305. Available from: [https://doi.org/10.1016/0022-247X\(78\)90181-6](https://doi.org/10.1016/0022-247X(78)90181-6).
- [7] Ali EE, Bulboaca T, Kota WY, El-Ashwah RM, Albalah AM. A study of generalized distribution series and their mapping properties in univalent function theory. *Aims Mathematics*. 2025; 10(6): 13296-13318. Available from: <https://www.aimspress.com/article/doi/10.3934/math.2025596>.
- [8] Attiya A, Ibrahim RW, Albalahi AM, Ali EE, Bulboaca T. A differential operator associated with q -raina function. *Symmetry*. 2022; 14: 1518. Available from: <https://doi.org/10.3390/sym14081518>.
- [9] Ali EE, Srivastava HM, Kota WY, El-Ashwah RM, Albalah AM. The second Hankel determinant and the Fekete-Szeg functional for a subclass of analytic functions by using the q -Salagean derivative operator. *Alexandria Engineering Journal*. 2025; 116: 141-146. Available from: <https://doi.org/10.1016/j.aej.2024.12.019>.
- [10] Esther OD, Olubunmi AF, Matthew OO. Results on multivalent Bessel functions associated with a new integral operator. *Partial Differential Equations in Applied Mathematics*. 2025; 14: 101228. Available from: <https://doi.org/10.1016/j.padiff.2025.101228>.
- [11] Matthew OO, Moses OB, Davids EO. On geometric properties of certain class of analytic functions connected with a multiplier transformation. *International Journal of Development Mathematics*. 2025; 2(2): 190-201. Available from: <https://orcid.org/0000-0003-3170-6818>.
- [12] Ali EE, El-Ashwah RM, Kota WY, Albalah AM. Geometric attributes of analytic functions generated by mittagleffler function. *Mathematics*. 2025; 13(20): 3284. Available from: <https://doi.org/10.3390/math13203284>.
- [13] Saleem N, Shafqat R, George R, Hussain A, Yassen M. A theoretical analysis on the fractional fuzzy controlled evolution equation. *Fractals*. 2023; (31)10: 107299. Available from: <https://doi.org/10.1142/S0218348X2340090X>.
- [14] Abdou Amir FI, Moussaoui A, Shafqat R, El Omari M, Melliani S. The Hadamard ψ -Caputo tempered fractional derivative in various types of fuzzy fractional differential equations. *Soft Computing*. 2024; 28: 9253-9270. Available from: <https://doi.org/10.1007/s00500-024-09821-w>.
- [15] Ali EE, El-Ashwah RM, Albalah AM, Mohammed WW. Properties for close-to-convex and quasi-convex functions using q -linear operator. *Mathematics*. 2025; 13: 900. Available from: <https://doi.org/10.3390/math13060900>.
- [16] Ali EE, Lashin A, Albalahi AM. Coefficient estimates for some classes of bi-univalent function associated with Jackson q -difference operator. *Journal of Function Spaces*. 2022; 2022(1): 1-8. Available from: <https://doi.org/10.1155/2022/2365918>.
- [17] Miller SS, Mocanu PT. Differential subordinations and univalent functions. *Michigan Mathematical Journal*. 1981; 28: 157-171. Available from: <https://doi.org/10.1307/mmj/1029002507>.
- [18] Miller SS, Mocanu PT. *Differential Subordinations: Theory and Applications; Series on Monographs and Textbooks in Pure and Applied Mathematics*. New York: Marcel Dekker; 2000. Available from: <https://doi.org/10.1201/9781482289817>.
- [19] Bulboaca T. *Differential Subordinations and Superordinations, Recent Results*. Romania: House of Scientific Book Publication; 2005.
- [20] Duren PL. *Univalent Functions*. Heidelberg: Springer; 1980.
- [21] Tayyah AS, Atshan WG. New results on r , k , μ -riemann-liouville fractional operators in complex domain with applications. *Fractal and Fractional*. 2024; 8(3): 165. Available from: <https://doi.org/10.3390/fractalfract8030165>.
- [22] Srivastava HM, Choi J. *Series Associated with the Zeta and Related Functions*. Netherlands: Springer Dordrecht; 2001.
- [23] Choi J, Srivastava HM. Certain families of series associated with the Hurwitz-Lerch Zeta functions. *Applied Mathematics and Computation*. 2005; 170(1): 399-409. Available from: <https://doi.org/10.1016/j.amc.2004.12.004>.

- [24] Ferreira C, Lopez JL. Asymptotic expansions of the Hurwitz-Lerch Zeta function. *Journal of Mathematical Analysis and Applications*. 2004; 298(1): 210-224. Available from: <https://doi.org/10.1016/j.jmaa.2004.05.040>.
- [25] Lin SD, Srivastava HM. Some families of the Hurwitz-Lerch Zeta functions and associated fractional derivatives and other integral representations. *Applied Mathematics and Computation*. 2004; 154(3): 725-733. Available from: [https://doi.org/10.1016/S0096-3003\(03\)00746-X](https://doi.org/10.1016/S0096-3003(03)00746-X).
- [26] El-Ashwah RM, Bulboacă T. Sandwich results for p -valent meromorphic functions associated with Hurwitz-Lerch Zeta function. *Tamsui Oxford Journal of Information and Mathematical Sciences*. 2018; 32(2): 48-63.
- [27] Atshan WG, Ali AR. On sandwich theorems results for certain univalent functions defined by generalized operators. *Iraqi Journal of Science*. 2021; 62(7): 2376-2383. Available from: <https://doi.org/10.24996/ij.s.2021.62.7.27>.
- [28] Shexo A, Atshan WG, Salim AJ. 3rd-order differential subordination and superordination results for univalent analytic functions associated with zeta-riemann fractional differential operator. *Journal of Al-Qadisiyah for Computer Science and Mathematics*. 2024; 16(3): 1-25. Available from: <https://doi.org/10.29304/jqcs.2024.16.31649>.
- [29] Eenigenburg P, Miller SS, Mocanu PT, Reade MO. On a Briot-Bouquet differential subordination. In: *General Inequalities 3*. Basel: Birkhäuser; 1983. p.339-348. Available from: https://doi.org/10.1007/978-3-0348-6290-5_26.
- [30] Padmanabhan KS, Parvathem R. Some applications of differential subordination. *Bulletin of the Australian Mathematical Society*. 1985; 32(3): 321-330. Available from: <https://doi.org/10.1017/S0004972700002410>.
- [31] Robertson RJ. On the theory of univalent functions. *Annals of Mathematics*. 1936; 37(2): 374-408. Available from: <https://doi.org/10.2307/1968451>.
- [32] Janowski W. Some extremal problems for certain families of analytic functions. *Annals of the Polish Mathematical Society*. 1973; 28(3): 297-326. Available from: <http://eudml.org/doc/266218>.
- [33] Libera RJ. Some classes of regular univalent functions. *Proceedings of the American Mathematical Society*. 1965; 16(4): 755-758. Available from: <https://doi.org/10.2307/2033917>.
- [34] Goel RM, Mahrok BS. Some invariance properties of a subclass of close-to-convex functions. *Indian Journal of Pure and Applied Mathematics*. 1981; 12(9): 1240-1249.
- [35] Kaplan W. Close-to-convex schlicht functions. *Michigan Mathematical Journal*. 1952; 1: 169-185.
- [36] Silvia EM. Subclasses of close-to-convex functions. *International Journal of Mathematics and Mathematical Sciences*. 1983; 3: 449-458. Available from: <https://doi.org/10.1155/S0161171283000393>.
- [37] Shafiq M, Khan N, Srivastava HM, Khan B, Ahmad QZ, Tahir M. Generalisation of close-to-convex functions associated with Janowski functions. *Maejo International Journal of Science and Technology*. 2020; 14(2): 141-155.
- [38] Chichra PN. New subclasses of the class of close-to-convex functions. *Proceedings of the American Mathematical Society*. 1977; 62: 37-43. Available from: <https://doi.org/10.1090/S0002-9939-1977-0425097-1>.
- [39] Zmorovich VA, Pokhilevich VA. On close-to-convex functions. *Ukrainian Mathematical Journal*. 1981; 33: 670-673.
- [40] Al-Amiri SH. Certain analogy of the-convex function. *Romanian Journal of Pure and Applied Mathematics*. 1978; 23: 1449-1454.
- [41] Miller SS, Mocanu PT, Reade MO. All α -convex functions are univalent and starlike. *Proceedings of the American Mathematical Society*. 1973; 37: 553-554. Available from: <https://doi.org/10.1090/S0002-9939-1973-0313490-3>.