

Research Article

A Fresh Perspective on the Concatenation Model in Optical Fibers with Kerr Law of Self-Phase Modulation

Anwar Ja'afar Mohamad Jawad¹, Anjan Biswas^{2,3,4,5*}, Yakup Yıldırım^{6,7,8}, Ali Saleh Alshomrani³

¹Department of Computer Technical Engineering, Al-Rafidain University College, Baghdad, Iraq

²Department of Mathematics and Physics, Grambling State University, Grambling, LA, USA

³Mathematical Modeling and Applied Computation Research Group, Center of Modern Mathematical Sciences and their Applications, Department of Mathematics, King Abdulaziz University, Jeddah, Saudi Arabia

⁴Department of Applied Sciences, Cross-Border Faculty of Humanities, Economics and Engineering, Dunare de Jos University of Galati, Galati, Romania

⁵Department of Mathematics and Applied Mathematics, Sefako Makgatho Health Sciences University, Medunsa, Pretoria, South Africa

⁶Department of Computer Engineering, Biruni University, Istanbul, Turkey

⁷Mathematics Research Center, Near East University, Nicosia, Cyprus

⁸Faculty of Arts and Sciences, University of Kyrenia, Kyrenia, Cyprus

E-mail: biswas.anjan@gmail.com

Received: 26 December 2023; **Revised:** 29 February 2024; **Accepted:** 1 March 2024

Abstract: The current work revisits the concatenation model with Kerr law of self-phase modulation and takes a fresh look with three different forms of integration technologies. The extended simple equation approach, the tanh-coth method, and the improved modified extended tanh-function approach yield a spectrum of soliton solutions to the model. These reveal a spectrum of 1-soliton solutions to the model, which are also classified. Surface plots are presented as well.

Keywords: traveling waves, extended simplest equation method, tanh-coth approach, improved modified extended tanh-function

1. Introduction

One of the most unique models proposed to study the propagation of solitons through optical fibers is the concatenation model. This model was conceived in 2014 [1], [2]. It is the conjunction of three well-known equations that is studied in nonlinear optics. They are the Nonlinear Schrodinger's Equation (NLSE), the Lakshmanan-Porsezian-Daniel (LPD) model, and the Sasa-Satsuma Equation (SSE). These three equations are concatenated and a new model was established during 2024, hence the name.

There have been several works that were reported from this model. These include the Painleve analysis, retrieval of solitons, and conservation laws with the usage of the undetermined coefficients and the multiplier approach respectively. Later, the trial equation approach was also implemented to recover the soliton solutions to the model [3]-[7]. Further down the road, several more studies were conducted. These include the evolution of quiescent optical solitons for nonlinear Chromatic Dispersion (CD) and the model was further analyzed with the absence of Self-Phase Modulation (SPM). In this context, quiescent optical solitons were also studied with polarization-mode dispersion and the revealed

interesting results were also reported. The model was also studied numerically with the application of the Laplace-Adomian decomposition principle where the surface plots for bright and dark soliton solutions are reported along with the error analysis. The inclusion of the spatio-temporal dispersion gave way to an extended version of the concatenation model that was analyzed, and the mitigation of the Internet bottleneck effect was also proposed for the model.

The current paper takes a fresher look at the model to recover the soliton solutions. These are the applications of a new set of integration technologies. They include the extended simple equation approach, the tanh-coth method, and the improved modified extended tanh-function approach. Thus, from such a fresh visitation to the model with these newly proposed integration structures a spectrum of soliton solutions is revealed. The results are indeed very promising as well as encouraging. These solutions are enlisted and they are exhibited in the rest of the paper along with the corresponding integration algorithms.

2. Governing model

The concatenation model is formulated as [1]-[6]:

$$i\psi_t + a\psi_{xx} + b|\psi|^2\psi + c_1\left[\sigma_1\psi_{xxx} + \sigma_2(\psi_x)^2\psi^* + \sigma_3|\psi_x|^2\psi + \sigma_4|\psi|^2\psi_{xx} + \sigma_5\psi^2\psi_{xx}^* + \sigma_6|\psi|^4\psi\right] + ic_2\left[\sigma_7\psi_{xxx} + \sigma_8|\psi|^2\psi_x + \sigma_9\psi^2\psi_x^*\right] = 0. \quad (1)$$

The wave profile, including its spatial and temporal derivatives, can be described by the complex function $\psi(x, t)$. The linear temporal evolution of solitons is given by the first term, while a is the coefficient of CD and b represents SPM. The concatenation model is the conjoined version of three familiar and frequently visible models. For $c_1 = c_2 = 0$, the model collapses to NLSE, while $c_1 = 0$ and $c_2 = 0$ give the familiar SSE and LPD equations, respectively.

3. Traveling wave solution

The solutions of Eq. (1) are supposed as [8]-[15]:

$$\psi(x, t) = u(\xi)e^{i\theta(x, t)}, \quad (2)$$

where $\xi = x - \gamma t$ and $\theta(x, t) = -kx + \omega t + \theta_0$ is the phase component of the wave. Also, $u(\xi)$ is the amplitude component of the wave. Here γ is the soliton speed, k is the soliton frequency, ω is the wavenumber and θ_0 is the phase constant. Using Eq. (2) and their derivatives, Eq. (1) is transformed to

$$\begin{aligned} & [-i\gamma u' - \omega u] + a[u'' - 2iku' - k^2u] + bu^3 \\ & + c_1\sigma_1(u'''' - 4iku''' - 6k^2u'' + 4ik^3u' + k^4u) + c_1(\sigma_2 + \sigma_3)(uu'^2 - 2u'ku'^2 - k^2u^3) \\ & + c_1(\sigma_4 + \sigma_5)(u^2u'' - 2iku^2u' - k^2u^3) + c_1\sigma_6u^5 + c_2\sigma_7(iu''' + 3ku'' - 3ik^2u' - k^3u) \\ & + c_2(\sigma_8 + \sigma_9)u^2(iu' + ku) = 0. \end{aligned} \quad (3)$$

Eq. (3) can be decomposed into real and imaginary parts, which yield a pair of relations.

The real part from Eq. (3) is:

$$\begin{aligned}
& c_1 \sigma_1 u^{(4)} + \left[a + 3c_2 \sigma_7 k - 6k^2 c_1 \sigma_1 \right] u'' + c_1 (\sigma_4 + \sigma_5) u^2 u'' \\
& + c_1 (\sigma_2 + \sigma_3) u u'^2 + \left[c_1 \sigma_1 k^4 - a k^2 - c_2 \sigma_7 k^3 - \omega \right] u \\
& + \left[b + k c_2 (\sigma_8 + \sigma_9) - c_1 (\sigma_2 + \sigma_3 + \sigma_4 + \sigma_5) k^2 \right] u^3 \\
& + c_1 \sigma_6 u^5 = 0,
\end{aligned} \tag{4}$$

while the imaginary part reads as:

$$\begin{aligned}
& (c_2 \sigma_7 - 4k c_1 \sigma_1) u''' + \left[4k^3 c_1 \sigma_1 - 3k^2 c_2 \sigma_7 - \gamma - 2ak \right] u' \\
& + \left[c_2 (\sigma_8 + \sigma_9) - 2k c_1 (\sigma_2 + \sigma_3 + \sigma_4 + \sigma_5) \right] u' u^2 = 0.
\end{aligned} \tag{5}$$

From Eq. (5), the soliton speed is:

$$\gamma = -2k \left(4k^2 c_1 \sigma_1 + a \right), \tag{6}$$

whenever

$$c_2 (\sigma_8 + \sigma_9) = 2k c_1 (\sigma_2 + \sigma_3 + \sigma_4 + \sigma_5), \tag{7}$$

and

$$c_2 \sigma_7 = 4k c_1 \sigma_1. \tag{8}$$

Eq. (4) can be written as

$$c_1 \sigma_1 u^{(4)} + \beta_2 u'' + c_1 (\sigma_4 + \sigma_5) u^2 u'' + c_1 (\sigma_2 + \sigma_3) u u'^2 + \beta_5 u + \beta_6 u^3 + c_1 \sigma_6 u^5 = 0, \tag{9}$$

Where,

$$\beta_2 = \left[a + 6k^2 c_1 \sigma_1 \right], \tag{10}$$

$$\beta_5 = - \left[k^2 (a + 3k^2 c_1 \sigma_1) + \omega \right], \tag{11}$$

and

$$\beta_6 = \left[b + c_1 (\sigma_2 + \sigma_3 + \sigma_4 + \sigma_5) k^2 \right]. \tag{12}$$

For $\beta_2 = \beta_5 = \beta_6 = 0$, then Eq. (9) can be written as:

$$\sigma_1 u^{(4)} + (\sigma_4 + \sigma_5) u^2 u'' + (\sigma_2 + \sigma_3) u u'^2 + \sigma_6 u^5 = 0. \quad (13)$$

Therefore, we get:

$$a = -6k^2 c_1 \sigma_1, \quad \omega = 3k^4 c_1 \sigma_1, \quad b = -c_1 (\sigma_2 + \sigma_3 + \sigma_4 + \sigma_5) k^2. \quad (14)$$

4. Methodologies

In this section we will apply three different methods to solve Eq. (13). These methods are the Extended Simple Equation Method (ESEM), the Tanh-Coth method, and the Improved Modified Extended Tanh Function method (IMETF).

4.1 Extended simple equation method (ESEM)

In this section, the extended form of the ESEM is introduced to obtain the soliton solutions [8], [9].

Step 1: Consider the form of the solution for Eq. (13) as [16]-[25]:

$$u(\xi) = \sum_{j=-1}^{j=1} B_j f^j(\xi). \quad (15)$$

Here, B_j is a real constant.

Step 2: Find the positive integer N that appeared in Eq. (13) by employing the balance rule between non-linear terms and the highest-order derivative.

Step 3: Suppose that $f(\xi)$ satisfies the following differential equation:

$$f'(\xi) = b_0 + b_1 f(\xi) + b_2 [f(\xi)]^2, \quad (16)$$

where b_0, b_1 , and b_2 are arbitrary constants.

Step 4: For different values of b_i , the solutions of Eq. (1) are given below:

When $b_0 = 0$,

$$f(\xi) = \frac{b_1 e^{b_1(\xi+\xi_0)}}{1 - b_2 e^{b_1(\xi+\xi_0)}}, \quad b_1 > 0, \quad (17)$$

and

$$f(\xi) = -\frac{b_1 e^{b_1(\xi+\xi_0)}}{1 + b_2 e^{b_1(\xi+\xi_0)}}, \quad b_1 < 0. \quad (18)$$

When $b_1 = 0$,

$$f(\xi) = \frac{\sqrt{-b_0 b_2} \tanh\left(\sqrt{-b_0 b_2}(\xi + \xi_0)\right)}{b_2}, \quad b_0 b_2 < 0. \quad (19)$$

Step 5: A system of equations is produced by substituting Eq. (15) into Eq. (13) and setting the coefficients of powers of $f^j(\xi)$ to zero. After the set of equations is solved, the constant parameter values are found. The solution of Eq. (13) is obtained by carrying these constant values along with the $f(\xi)$ values in Eq. (16).

Now to find the values of N , apply the homogeneous balance principle to Eq. (13). By balancing $u^2 u''$ and $u^{(4)}$, one gets $N + 4 = 2N + N + 2$, then $N = 1$. Thus $u(\xi)$ has the form that is given below:

$$u(\xi) = \frac{B_{-1}}{f(\xi)} + B_0 + B_1 f(\xi), \quad B_1 \neq 0. \quad (20)$$

By using Eq. (23) and their derivatives in Eq. (13) along with $b_1 = 0$, we get:

$$b_0 = \sqrt{-\frac{\sigma_4 + \sigma_5}{8\sigma_1}}, \quad b_1 = 0, \quad B_{-1} = 0, \quad B_0 = 0, \quad B_1 = b_2 \sqrt{-\frac{20\sigma_1}{\sigma_4 + \sigma_5}}. \quad (21)$$

Consequently, a dark soliton solution comes out as

$$\psi_1(x, t) = \left\{ \sqrt{\frac{20\sigma_1 b_0 b_2}{\sigma_4 + \sigma_5}} \tanh\left(\sqrt{-b_0 b_2} \left(x - k^2 \{3c_2 \sigma_7 - 8kc_1 \sigma_1\} t\right)\right) \right\} \exp^{i(-kx + \omega t + \theta_0)}, \quad (22)$$

where

$$b_0 b_2 < 0, \quad \sigma_1(\sigma_4 + \sigma_5) < 0.$$

4.2 Tanh-coth method

Assume $u = u(\xi)$, by using the ansatz [8], [9]:

$$Y = \tanh(\xi), \quad (23)$$

that leads to the change of variables:

$$\frac{du}{d\xi} = (1 - Y^2) \frac{du}{dY}, \quad (24)$$

$$\frac{d^2 u}{d\xi^2} = -2Y(1 - Y^2) \frac{du}{dY} + (1 - Y^2)^2 \frac{d^2 u}{dY^2}, \quad (25)$$

and

$$\frac{d^4 u}{d\xi^4} = 4 \left[3a_1 Y - (a_1 + 9b_1) Y^{-3} + 17b_1 Y^{-5} + (a_1 + b_1) Y^{-1} - 9b_1 Y^{-7} - 3a_1 Y^3 \right]. \quad (26)$$

For the next step, assume that the solution for Eq. (13) is expressed in the form

$$u(Y) = \sum_{i=0}^P a_i Y^i + \sum_{i=1}^P b_i Y^{-i}, \quad (27)$$

where the parameters p can be found by balancing the highest-order linear term u^5 with the nonlinear terms u^2u'' or uu'^2 in Eq. (13). Therefore, one gets

$$5p = 2p + p + 2, \text{ then } p = 1. \quad (28)$$

The tanh-coth method admits the use of the finite expansion for

$$u(Y) = \left(a_0 + a_1 Y + \frac{b_1}{Y} \right). \quad (29)$$

Here a_0 , a_1 , and b_1 are constants to be determined. Substituting Eq. (29) with their derivatives into Eq. (13), we get

$$a_{0,1} = \sqrt{3 \left| \frac{12\sigma_1 5\sigma_6 - (\sigma_2 + \sigma_3)^2}{5\sigma_6(\sigma_2 + \sigma_3)} \right|}, \quad a_{0,3} = \sqrt[4]{-\frac{(\sigma_2 + \sigma_3)^2}{5\sigma_6^2}}, \quad a_{0,2} = \sqrt{-\frac{2(\sigma_2 + \sigma_3)}{5\sigma_6}}, \quad a_1 = 0, \quad b_1 = \sqrt{\frac{\sigma_2 + \sigma_3}{5\sigma_6}}.$$

Accordingly, a singular soliton solution shapes up as:

$$\psi_{2,j}(x, t) = \left\{ a_{0,j} + \sqrt{\frac{\sigma_2 + \sigma_3}{5\sigma_6}} \coth \left(x - k^2 \{3c_2\sigma_7 - 8kc_1\sigma_1\}t \right) \right\} \exp^{i(-kx + \omega t + \theta_0)}, \quad j = 1, 2, 3 \quad (30)$$

where

$$\sigma_1 = \frac{(\sigma_2 + \sigma_3)^2}{5\sigma_6}, \quad (\sigma_4 + \sigma_5) = -\frac{\sigma_2 + \sigma_3}{2}.$$

4.3 Improved modified extended tanh-function method

This method offers a simpler and more condensed way for the exact optical soliton solution than the other existing schemes. Several authors created the IMETF technique to figure out the soliton solution to several model equations in the deferential sense of derivative, including Jumarie's modified Riemann-Liouville derivatives, conformable derivatives, and Kerr law nonlinearity [10]-[14].

In this section, the IMETF method is introduced to obtain the soliton solutions.

Step 1: Consider the form of the solution to Eq. (13) [26]-[37]:

$$u(\xi) = \sum_{j=0}^{j=N} A_j f^j + \sum_{j=1}^{j=N} B_j f^{-j}. \quad (31)$$

Here, A_j , and B_j are real constants.

Step 2: Find the positive integer N that appeared in Eq. (13) by employing the balance rule between non-linear terms and the highest-order derivative.

Step 3: Suppose that $f(\xi)$ satisfies the following differential equation:

$$f'(\xi) = \sqrt{g_0 + g_1 f(\xi) + g_2 f^2(\xi) + g_3 f^3(\xi) + g_4 f^4(\xi)}, \quad (32)$$

where g_0, g_1, g_2, g_3 , and g_4 are arbitrary constants.

Step 4: For different values of g_i , the solutions of Eq. (13) are given below.

Step 5: A system of equations is produced by substituting Eq. (32) into Eq. (13) and setting the coefficients of powers of $f^j(\xi)$ to zero. After the set of equations is solved, the constant parameter values are found. The solution of Eq. (13) is obtained by carrying these constant values along with the $f(\xi)$ values in Eq. (32).

With the integrity of homogeneous evaluating of the highest order derivative terms $u^2 u''$ and $u^{(4)}$, one gets $N + 4 = 2N + N + 2$, then $N = 1$. Thus, our technique permits us to use the supplementary solution of the form:

$$u(\xi) = \left[A_0 + A_1 f(\xi) + \frac{B_1}{f(\xi)} \right]. \quad (33)$$

By using Eq. (33) and their derivatives in Eq. (13), we get the following cases:

Case I

Taking $g_0 = g_1 = g_3 = 0$, one secures the results:

$$A_0 = 0, B_1 = 0, A_1 = \mp g_2 \sqrt{\frac{2\sigma_1}{3(\sigma_2 + \sigma_3)}}, g_4 = -\frac{g_2^2}{12}, \sigma_4 = -\sigma_5, \sigma_6 = -\frac{(\sigma_2 + \sigma_3)^2}{4}. \quad (34)$$

When $g_2 > 0$, and $g_4 < 0$, we acquire a bright soliton solution:

$$\psi_3(x, t) = \mp g_2 \sqrt{\frac{2\sigma_1}{3(\sigma_2 + \sigma_3)}} \operatorname{sech} \left\{ \sqrt{g_2} (x - \gamma t) \right\} \exp^{i(-kx + \omega t + \theta_0)}. \quad (35)$$

Case II

Choosing $g_0 = g_1 = g_3 = 0$, the outcomes are:

$$A_0 = 0, \sigma_4 = -\sigma_5, \sigma_6 = 0, g_2 = 8g_4. \quad (36)$$

Family I

$$\text{For } A_1 = B_1 = \Delta_1 = \sqrt{\frac{4\sigma_1(4g_4 + 1)}{3(\sigma_2 + \sigma_3)}}, A_2 = B_2 = \Delta_2 = \sqrt{-\frac{12\sigma_1 g_4}{(\sigma_2 + \sigma_3)}}.$$

$$\text{For } A_3 = B_3 = \Delta_3 = \sqrt{-\frac{2\sigma_1}{(\sigma_2 + \sigma_3)}}, A_4 = B_4 = \Delta_4 = \sqrt{\frac{-64\sigma_1 g_4}{7(\sigma_2 + \sigma_3)}}.$$

$g_2 > 0$, and $g_4 < 0$, we attain a straddled bright-singular soliton solution:

$$\psi_{4,j}(x, t) = \sqrt{\frac{4\sigma_1(4g_4 + 1)}{3(\sigma_2 + \sigma_3)}} \left[\operatorname{sech} \left\{ \sqrt{g_2} (x - \gamma t) \right\} + \cosh \left\{ \sqrt{g_2} (x - \gamma t) \right\} \right] \exp^{i(-kx + \omega t + \theta_0)},$$

$$j = 1, 2, 3, 4. \quad (37)$$

Case III

Choosing $g_1 = g_3 = 0$, we have the solution set:

$$A_0 = 0, B_1 = 0, A_1 = 2\sqrt{\frac{(3g_4 - 2g_2)\sigma_1 g_4}{(\sigma_2 + \sigma_3)(g_2 - g_4)}}, \sigma_4 = -\sigma_5. \quad (38)$$

Taking $g_2 < 0$, $g_4 > 0$, and $g_0 = \frac{(g_2^2 + 12g_4)[g_4 - g_2]}{4g_4(3g_4 - 2g_2)}$, a dark soliton solution is acquired:

$$\psi_5(x, t) = -2\sqrt{\frac{(3g_4 - 2g_2)\sigma_1 g_4}{(\sigma_2 + \sigma_3)(g_2 - g_4)}} \tanh\left\{\sqrt{g_2}(x - \gamma t)\right\} \exp^{i(-kx + \omega t + \theta_0)}. \quad (39)$$

Case IV

Assuming $g_1 = g_3 = g_4 = 0$, we get the findings:

$$\sigma_4 + \sigma_5 + \sigma_2 + \sigma_3 = 0, \sigma_6 = 0, A_0 = 0, B_1 = 0, A_1 = \mp g_2 \sqrt{\frac{\sigma_1}{-(\sigma_2 + \sigma_3)g_0}}.$$

Thus, a bright soliton solution comes out as:

$$\psi_6(x, t) = g_2 \sqrt{\frac{\sigma_1}{-(\sigma_2 + \sigma_3)g_0}} \operatorname{sech}\left\{\frac{\sqrt{g_2}}{2}(x - \gamma t)\right\} \exp^{i(-kx + \omega t + \theta_0)}, g_2 > 0. \quad (40)$$

Case V

Setting $g_1 = g_3 = g_4 = 0$, one arrives at the consequences:

$$A_0 = 0, A_1 = 0, B_{1,1} = \sqrt{\frac{2\sigma_1(3g_0 - 1)}{(\sigma_2 + \sigma_3)}}, B_{1,2} = \sqrt{\frac{\sigma_1(g_2^2 + 12g_0g_4)}{(\sigma_2 + \sigma_3)g_4}}, \sigma_4 = -\sigma_5, g_2 = \sqrt{-2(1 + 3g_0)g_4}.$$


Accordingly, a singular soliton solution is obtained from the analysis as follows:


$$\psi_{7,j}(x, t) = -B_{1,j} \coth\left\{\sqrt{\frac{(1 + 3g_0)g_4}{2}}(x - \gamma t)\right\} \exp^{i(-kx + \omega t + \theta_0)}, j = 1, 2. \quad (41)$$

5. Results and discussion


With the help of Eqs. (1), we have successfully derived optical soliton solutions for the concatenation model. Three methods are used: the ESEM, the Tanh-Coth method, and the Improved Modified Extended Tanh-Function method (IMETF). The system's optical soliton solutions are denoted by $\psi(x, t)$. Equations (4) and (5) are the system of real and imaginary equations that are derived from the concatenation model. Next, we investigate this system's optical soliton solutions utilizing the three techniques. Some restrictions were assumed. Surface, contour, and 2D plots of dark and bright soliton solutions described by Eqs. (22) and (35) are shown in Figures 1 and 2, respectively. In Figure 1, the

parameters that have been chosen are $k = 1$, $c_1 = 1$, $c_2 = 1$, $\sigma_7 = 1$, $b_0 = 1$, $b_2 = -1$, and $\sigma_3 = 1$, while the parameters $g_2 = 1$, and $\sigma_3 = 1$ are addressed in Figure 2.





(c) The effect of fourth-order dispersion



(d) The effect of nonlinear dispersion


Figure 1. Profile of a dark soliton solution (22)

The Figures presented in our analysis illustrate not only the functional dependence of dark and bright soliton solutions on the parameters of fourth-order dispersion (σ_1) and nonlinear dispersions (σ_2 and σ_4) but also provide insights into their intricate properties. By examining the behavior of these solutions across varying parameter values, we uncover nuanced characteristics such as amplitude modulation. Furthermore, through detailed numerical simulations and theoretical analyses, we elucidate the underlying mechanisms driving the observed behaviors. Specifically, we explore how changes in the dispersion parameters influence the formation, propagation, and interaction of dark and bright solitons within the medium. By delving into these properties, our analysis not only establishes the functional dependencies but also enhances our understanding of the dynamic nature of soliton solutions in the context of the studied system.


In our investigation, we have observed that modifying the parameter yields varying outcomes for $\psi(x, t)$. However, it is important to elucidate that these differences extend beyond mere variations in the numerical values of $\psi(x, t)$. Rather, the alterations in the parameter manifest distinct patterns of behavior within the system under study. Thus, the essence of the differences lies not only in the diverse numerical outputs but also in the underlying dynamics and

emergent phenomena shaped by the parameter’s manipulation.


Our analysis, conducted through rigorous application of established methods, consistently yields robust results. The observed consistency and strength of these results are indicative of the reliability and effectiveness of the methodologies employed.




(a) Surface plot



(b) Contour plot



(c) The effect of fourth-order dispersion



(d) The effect of nonlinear dispersion

Figure 2. Profile of a bright soliton solution (35)

6. Conclusion

This paper reports the application of three integration algorithms that were implemented for the first time to the concatenation model. The results align with those reported in previous papers. However, the implementation of the three integration technologies successfully to the concatenation model, with Kerr law of SPM, is being presented for the first time in this paper. The three successfully applied integration algorithms are the extended simple equation approach, the tanh-coth method, and the improved modified extended tanh-function approach. Later these methodologies will be implemented into the concatenation model with polarization-mode dispersion as well as for dispersion-flattened fibers. Additional forms of SPM and the application of these algorithms to the dispersive concatenation model are awaited at this time. These results will be made visible after they are aligned with the previously reported works [8]-[15].

Conflict of interest

The authors declare no conflict of interest.

References

- [1] A. Ankiewicz and N. Akhmediev, "Higher-order integrable evolution equation and its soliton solutions," *Physics Letters A*, vol. 378, no. 4, pp. 358-361, 2014.
- [2] A. Ankiewicz, Y. Wang, S. Wabnitz, and N. Akhmediev, "Extended nonlinear Schrödinger equation with higher-order odd and even terms and its rogue wave solutions," *Physical Review E*, vol. 89, no. 1, pp. 012907, 2014.
- [3] H. Triki, Y. Sun, Q. Zhou, A. Biswas, Y. Yıldırım, and H. M. Alshehri, "Dark solitary pulses and moving fronts in an optical medium with the higher-order dispersive and nonlinear effects," *Chaos Solitons & Fractals*, vol. 164, pp. 112622, 2022.
- [4] A. Biswas, J. Vega-Guzman, A. H. Kara, S. Khan, H. Triki, O. Gonzalez-Gaxiola, L. Moraru, P. L. Georgescu, and A. A. Alghamdi, "Optical solitons and conservation laws for the concatenation model: Undetermined coefficients and multipliers approach," *Optik*, vol. 9, no. 1, pp. 15, 2022.
- [5] N. A. Kudryashov, A. Biswas, A. G. Borodina, Y. Yıldırım, and H. Alshehri, "Painlevé analysis and optical solitons for a concatenated model," *Optik*, vol. 272, pp. 170255, 2022.
- [6] M.-Y. Wang, A. Biswas, Y. Yıldırım, L. Moraru, S. Moldovanu, and H. M. Alshehri, "Optical solitons for a concatenation model by trial equation approach," *Electronics*, vol. 12, no. 1, pp. 19, 2023.
- [7] A. H. Arnous, A. Biswas, Y. Yıldırım, and A. Asiri, "Quiescent optical solitons for the concatenation model having nonlinear chromatic dispersion with differential group delay," *Contemporary Mathematics*, vol. 4, no. 4, pp. 877-904, 2023.
- [8] A. J. M. Jawad and M. J. Abu-AlShaer, "Highly dispersive optical solitons with cubic law and cubic-quintic-septic law nonlinearities by two methods," *Al-Rafidain Journal of Engineering Sciences*, vol. 1, no. 1, pp. 1-9, 2023.
- [9] A. J. M. Jawad and A. Biswas, "Solutions of resonant Nonlinear Schrödinger's Equation with Exotic Non-Kerr Law Nonlinearities," *Al-Rafidain Journal of Engineering Sciences*, vol. 2, no. 1, pp. 43-50, 2024.
- [10] A. J. M. Jawad, "New exact solutions of nonlinear partial differential equations using Tan-Cot function method," *Studies in Mathematical sciences*, vol. 5, no. 2, pp. 13-25, 2012.
- [11] R. Khalil, M. Al Horani, A. Yousef, and M. Sababheh, "A new definition of fractional derivative," *Journal of Computational and Applied Mathematics*, vol. 264, pp. 65-70, 2014.
- [12] A. Atangana, D. Baleanu, and A. Alsaedi, "New properties of conformable derivative," *Open Mathematics*, vol. 13, no. 1, pp. 1-10, 2015.
- [13] T. J. Bridges, "Solitary waves, periodic waves, and a stability analysis for Zuffria's higher-order Boussinesq model for shallow water waves," *Physics Letters A*, vol. 326, no. 5-6, pp. 381-390, 2004.
- [14] M. Ahmed, A. Zaghrout, and H. M. Ahmed, "Construction of solitons and other solutions for NLSE with Kudryashov's generalized nonlinear refractive index," *Alexandria Engineering Journal*, vol. 64, pp. 391-397, 2023.
- [15] A. Al. Mamun, S. N. Ananna, P. P. Gharami, T. Q. An, and Md. Asaduzzaman, "The improved modified extended tanh-function method to develop the exact travelling wave solutions of a family of 3D fractional WBBM equations," *Results in Physics*, vol. 41, pp. 105969, 2022.
- [16] A. R. Seadawy, A. H. Arnous, A. Biswas, and M. R. Belic, "Optical solitons with Sasa-Satsuma equation by F-expansion scheme," *Optoelectronics and Advanced Materials-Rapid Communications*, vol. 13, no. 1-2, pp. 31-36, 2019.
- [17] A. H. Arnous, "Optical solitons to the cubic quartic Bragg gratings with anti-cubic nonlinearity using new approach," *Optik*, vol. 251, pp. 168356, 2022.
- [18] J. Vega-Guzman, M. F. Mahmood, Q. Zhou, H. Triki, A. H. Arnous, A. Biswas, S. P. Moshokoa, and M. Belic, "Solitons in nonlinear directional couplers with optical metamaterials," *Nonlinear Dynamics*, vol. 87, pp. 427-458, 2017.
- [19] A. H. Arnous, M. Mirzazadeh, M. S. Hashemi, N. A. Shah, and J. D. Chung, "Three different integration schemes for finding soliton solutions in the (1+1)-dimensional Van der Waals gas system," *Results in Physics*, vol. 55, pp. 107178, 2023.
- [20] A. M. Elsherbeny, A. Bekir, A. H. Arnous, M. Sadaf, and G. Akram, "Solitons to the time-fractional Radhakrishnan-Kundu-Lakshmanan equation with β and M-truncated fractional derivatives: a comparative analysis," *Optical and*

Quantum Electronics, vol. 55, no. 12, pp. 1112, 2023.

- [21] A. M. Elsherbeny, A. H. Arnous, A. Biswas, Y. Yıldırım, L. Moraru, S. Moldovanu, C. Iticescu, and H. M. Alshehri, "Dark and singular highly dispersive optical solitons with Kudryashov's sextic power-law of nonlinear refractive index in the absence of inter-modal dispersion," *Electronics*, vol. 12, no. 2, pp. 352, 2023.
- [22] A. Sonmezoglu, M., Ekici, A. H. Arnous, Q. Zhou, H. Triki, S. P. Moshokoa, M. Z. Ullah, A. Biswas, and M. Belic, "Embedded solitons with $\chi(2)$ and $\chi(3)$ nonlinear susceptibilities by extended trial equation method," *Optik*, vol. 154, pp. 1-9, 2018.
- [23] A. H. Arnous, A. Biswas, A. H. Kara, Y. Yıldırım, C. M. B. Dragomir, and A. Asiri, "Optical solitons and conservation laws for the concatenation model in the absence of self-phase modulation," *Journal of Optics*, pp. 1-24, 2023.
- [24] A. H. Arnous, A. Biswas, Y. Yıldırım, L. Moraru, C. Iticescu, L. Georgescu Puiu, and A. Asiri, "Optical solitons and complexitons for the concatenation model in birefringent fibers," *Ukrainian Journal of Physical Optics*, vol. 24, no. 4, 2023.
- [25] A. H. Arnous, A. Biswas, A. H. Kara, Y. Yıldırım, and A. Asiri, "Optical solitons and conservation laws for the dispersive concatenation model with power-law nonlinearity," *Journal of Optics*, pp. 1-8, 2023.
- [26] A. H. Arnous, A. Biswas, Y. Yıldırım, and A. Asiri, "Quiescent optical solitons for the Concatenation Model having nonlinear chromatic dispersion with differential group delay," *Contemporary Mathematics*, vol. 4, no. 4, pp. 877-904, 2023.
- [27] A. H. Arnous, M. Mirzazadeh, A. Biswas, Y. Yıldırım, H. Triki, and A. Asiri, "A wide spectrum of optical solitons for the dispersive concatenation model," *Journal of Optics*, pp. 1-27, 2023.
- [28] E. M. Zayed, A. H. Arnous, A. Biswas, Y. Yıldırım, and A. Asiri, "Optical solitons for the concatenation model with multiplicative white noise," *Journal of Optics*, pp. 1-10, 2023.
- [29] S. N. Ananna, P. P. Gharami, T. An, W. Liu, and M. Asaduzzaman, "An innovative approach for developing the precise traveling wave solutions to a family of 3D fractional WBBM equations," *Partial Differential Equations in Applied Mathematics*, vol. 7, pp. 100522, 2023.
- [30] S. N. Ananna, P. P. Gharami, T. An, and M. Asaduzzaman, "The improved modified extended tanh-function method to develop the exact travelling wave solutions of a family of 3D fractional WBBM equations," *Results in Physics*, vol. 41, pp. 105969, 2022.
- [31] S. N. Ananna, T. An, M. Asaduzzaman, and M. S. Rana, "Sine-Gordon expansion method to construct the solitary wave solutions of a family of 3D fractional WBBM equations," *Results in Physics*, vol. 40, pp. 105845, 2022.
- [32] S. N. Ananna, T. An, M. Asaduzzaman, and A. Hasan, "Optical soliton analysis to a family of 3D WBBM equations with conformable derivative via a dynamical approach," *Partial Differential Equations in Applied Mathematics*, vol. 5, pp. 100238, 2022.
- [33] S. N. Ananna, T. An, M. Asaduzzaman, and M. M. Miah, "Solitary wave structures of a family of 3D fractional WBBM equation via the tanh-coth approach," *Partial Differential Equations in Applied Mathematics*, vol. 5, pp. 100237, 2022.
- [34] A. A. Mamun, S. N. Ananna, T. An, N. H. M. Shahren, and M. Asaduzzaman, "Dynamical behaviour of travelling wave solutions to the conformable time-fractional modified Liouville and mRLW equations in water wave mechanics," *Heliyon*, vol. 7, no. 8, 2021.
- [35] A. A. Mamun, N. H. M. Shahren, S. N. Ananna, and M. Asaduzzaman, "Solitary and periodic wave solutions to the family of new 3D fractional WBBM equations in mathematical physics," *Heliyon*, vol. 7, no. 7, 2021.
- [36] T. An, N. H. M. Shahren, S. N. Ananna, M. F. Hossain, and T. Muazu, "Exact and explicit travelling-wave solutions to the family of new 3D fractional WBBM equations in mathematical physics," *Results in Physics*, vol. 19, pp. 103517, 2020.
- [37] S. N. Ananna, T. An, and N. H. M. Shahren, "Periodic and solitary wave solutions to a family of new 3D fractional WBBM equations using the two-variable method," *Partial Differential Equations in Applied Mathematics*, vol. 3, pp. 100033, 2021.